



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 7, 2021 – 10:38 am BST

PDB ID : 7BOC  
Title : Crystal structure of the PRMT5 TIM barrel domain in complex with RioK1 peptide  
Authors : Krzyzanowski, A.; t Hart, P.; Waldmann, H.; Gasper, R.  
Deposited on : 2021-01-25  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.23.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	 5% 48% 20% 28%
2	B	15	 20% 40% 40% 13% 7%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3266 atoms, of which 1422 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein arginine N-methyltransferase 5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	211	3163	1130	1422	304	298	9	0	0	0


- Molecule 2 is a protein called peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	14	103	61	18	24	0	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein arginine N-methyltransferase 5



- Molecule 2: peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.63Å 96.63Å 112.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.31 – 2.55 48.32 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.31-2.55) 99.9 (48.32-2.55)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 2.54Å)	Xtrriage
Refinement program	PHENIX 1.18-3855	Depositor
R, $R_{free}$	0.225 , 0.251 0.227 , 0.255	Depositor DCC
$R_{free}$ test set	1015 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.1	Xtrriage
Anisotropy	0.509	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 57.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3266	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	92.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.50	0/1791	0.74	2/2428 (0.1%)
2	B	0.44	0/104	0.59	0/140
All	All	0.49	0/1895	0.74	2/2568 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	LEU	CB-CG-CD1	-5.54	101.59	111.00
1	A	157	LEU	CB-CG-CD2	-5.51	101.64	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1741	1422	1739	50	0
2	B	103	0	92	9	0
All	All	1844	1422	1831	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:PHE:CD1	2:B:7:GLN:HG3	2.05	0.91
1:A:49:ARG:HD3	1:A:79:ASN:ND2	1.98	0.77
1:A:163:LEU:O	1:A:164:ARG:HG2	1.90	0.71
1:A:199:SER:HB3	1:A:202:ILE:HD12	1.74	0.69
1:A:189:TRP:CZ2	1:A:193:ARG:HD2	2.28	0.69
1:A:240:LYS:HE2	2:B:12:ASP:HB2	1.74	0.69
1:A:49:ARG:HD3	1:A:79:ASN:HD21	1.59	0.66
1:A:243:PHE:HD1	2:B:7:GLN:HG3	1.60	0.65
1:A:240:LYS:HE2	2:B:12:ASP:CB	2.28	0.64
1:A:254:ILE:HG22	1:A:258:LEU:HD11	1.79	0.63
1:A:45:VAL:HB	1:A:81:LEU:CD1	2.28	0.63
1:A:126:LEU:HD12	1:A:126:LEU:O	1.98	0.63
1:A:96:VAL:HG12	1:A:99:ILE:H	1.65	0.62
1:A:43:MET:HG3	1:A:44:PRO:HD2	1.83	0.59
1:A:50:PHE:CZ	1:A:285:GLU:HG3	2.38	0.59
1:A:190:HIS:O	1:A:194:THR:HG23	2.04	0.57
1:A:81:LEU:HD12	1:A:81:LEU:C	2.25	0.57
1:A:138:LEU:C	1:A:138:LEU:HD13	2.25	0.56
1:A:273:SER:OG	1:A:276:GLU:HG3	2.06	0.56
1:A:211:ASP:O	1:A:212:LEU:HD23	2.07	0.55
1:A:243:PHE:CE1	2:B:7:GLN:HG3	2.40	0.54
1:A:45:VAL:HB	1:A:81:LEU:HD12	1.90	0.54
1:A:138:LEU:HD13	1:A:138:LEU:O	2.09	0.53
1:A:248:LYS:NZ	2:B:10:ASP:OD1	2.38	0.53
1:A:238:THR:HA	1:A:243:PHE:O	2.10	0.51
1:A:90:ILE:HG23	1:A:103:SER:HB3	1.93	0.51
1:A:45:VAL:HB	1:A:81:LEU:HD11	1.92	0.50
1:A:92:PRO:HG3	1:A:107:MET:HE2	1.94	0.50
1:A:220:ARG:HG2	1:A:220:ARG:HH21	1.76	0.50
1:A:41:LEU:O	1:A:41:LEU:HD13	2.11	0.49
1:A:158:VAL:HG11	1:A:163:LEU:HD11	1.95	0.49
1:A:43:MET:HG3	1:A:44:PRO:CD	2.41	0.49
1:A:119:LEU:O	1:A:148:SER:HB2	2.13	0.48
1:A:254:ILE:O	1:A:258:LEU:HD12	2.14	0.47
1:A:96:VAL:O	1:A:100:ARG:HG3	2.14	0.47
1:A:163:LEU:HD13	1:A:211:ASP:CG	2.35	0.47
1:A:254:ILE:HG22	1:A:258:LEU:CD1	2.45	0.47
1:A:269:THR:HG22	1:A:269:THR:O	2.14	0.46
1:A:260:LEU:N	1:A:260:LEU:HD22	2.31	0.46
2:B:3:VAL:HA	2:B:7:GLN:OE1	2.15	0.46
1:A:189:TRP:NE1	1:A:224:GLU:OE1	2.38	0.46
1:A:220:ARG:HH21	1:A:220:ARG:CG	2.30	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:269:THR:O	1:A:272:HIS:HB2	2.18	0.43
1:A:247:SER:O	1:A:251:GLN:HG3	2.18	0.43
2:B:2:ARG:HD3	2:B:3:VAL:N	2.34	0.43
1:A:126:LEU:CD1	1:A:155:VAL:HG13	2.50	0.42
1:A:243:PHE:HB3	1:A:244:PRO:HD2	2.01	0.42
1:A:250:HIS:O	1:A:254:ILE:HG12	2.20	0.42
1:A:150:MET:HE1	1:A:200:LYS:O	2.20	0.41
1:A:91:ARG:HE	1:A:91:ARG:HB2	1.64	0.41
1:A:180:SER:O	1:A:180:SER:OG	2.36	0.41
1:A:214:SER:O	1:A:218:ILE:HG13	2.20	0.41
2:B:4:VAL:HB	2:B:5:PRO:HD2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/292 (70%)	194 (95%)	10 (5%)	1 (0%)	29	40
2	B	12/15 (80%)	10 (83%)	2 (17%)	0	100	100
All	All	217/307 (71%)	204 (94%)	12 (6%)	1 (0%)	29	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	214	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/254 (75%)	167 (87%)	24 (13%)	4	4
2	B	12/13 (92%)	9 (75%)	3 (25%)	0	0
All	All	203/267 (76%)	176 (87%)	27 (13%)	4	3

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	LEU
1	A	43	MET
1	A	47	HIS
1	A	50	PHE
1	A	51	LYS
1	A	96	VAL
1	A	101	ARG
1	A	109	GLN
1	A	127	ASN
1	A	149	SER
1	A	155	VAL
1	A	158	VAL
1	A	180	SER
1	A	204	VAL
1	A	214	SER
1	A	215	ASN
1	A	220	ARG
1	A	222	LEU
1	A	227	LYS
1	A	238	THR
1	A	249	MET
1	A	267	THR
1	A	271	HIS
1	A	275	LYS
2	B	1	SER
2	B	3	VAL
2	B	4	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	250	HIS
1	A	251	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	211/292 (72%)	0.65	15 (7%) 16 19	61, 82, 129, 171	0
2	B	14/15 (93%)	1.11	3 (21%) 0 0	83, 102, 128, 143	0
All	All	225/307 (73%)	0.68	18 (8%) 12 15	61, 83, 129, 171	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	163	LEU	5.5
1	A	179	TYR	5.3
1	A	164	ARG	4.4
2	B	1	SER	4.3
1	A	290	ASN	3.5
1	A	271	HIS	3.5
1	A	257	LEU	3.0
1	A	259	LYS	2.8
1	A	255	PHE	2.7
1	A	113	PHE	2.5
1	A	51	LYS	2.5
2	B	4	VAL	2.4
1	A	50	PHE	2.4
1	A	86	LEU	2.2
1	A	264	PHE	2.2
1	A	162	ASP	2.2
2	B	2	ARG	2.1
1	A	45	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.