Full wwPDB X-ray Structure Validation Report (i)

Apr 28, 2024 – 06:46 pm BST

PDB ID	:	4C1B
Title	:	Esterase domain of the ZfL2-1 ORF1 protein from the zebrafish ZfL2-1 retrotransposon
Authors	:	Schneider, A.M.; Weichenrieder, O.
Deposited on	:	2013-08-11
Resolution	:	2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	171	
1	B	171	
1	C	171	

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 8108 atoms, of which 4083 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ORF1-ENCODED PROTEIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	171	2740	853	1392	255	237	3	0	0	0
1	B	168	2709	843	1377	252	234	3	0	0	0
1	C	158	2585	807	1314	241	220	3	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	132	GLY	-	expression tag	UNP Q3LG57
A	133	PRO	-	expression tag	UNP Q3LG57
A	134	ALA	-	expression tag	UNP Q3LG57
A	135	MET	-	expression tag	UNP Q3LG57
B	132	GLY	-	expression tag	UNP Q3LG57
B	133	PRO	-	expression tag	UNP Q3LG57
B	134	ALA	-	expression tag	UNP Q3LG57
B	135	MET	-	expression tag	UNP Q3LG57
C	132	GLY	-	expression tag	UNP Q3LG57
C	133	PRO	-	expression tag	UNP Q3LG57
C	134	ALA	-	expression tag	UNP Q3LG57
C	135	MET	-	expression tag	UNP Q3LG57

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	38	38	38	0	0
2	B	25	25	25	0	0
2	C	11	11	11	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ORF1-ENCODED PROTEIN

Chain A:





- Molecule 1: ORF1-ENCODED PROTEIN

Chain B:





- Molecule 1: ORF1-ENCODED PROTEIN

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, α, β, γ	111.26Å 111.26Å 115.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.71 – 2.50 45.71 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.71-2.50) 100.0 (45.71-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	3.18 (at 2.51Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.167 , 0.210 0.169 , 0.211	Depositor DCC
R_{free} test set	1244 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å^2)	51.5	Xtriage
Anisotropy	0.202	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.41 , 46.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.006 for l,-k,h 0.016 for -l,-k,-h 0.013 for -h,-l,-k 0.004 for -h,l,k 0.034 for -h,k,-l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	8108	wwPDB-VP
Average B, all atoms (Å^2)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.53	0/1376	0.65	0/1865
1	B	0.55	0/1359	0.64	0/1841
1	C	0.49	0/1296	0.63	1/1754 (0.1%)
All	All	0.53	0/4031	0.64	1/5460 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	236	ARG	NE-CZ-NH2	-6.62	116.99	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1348	1392	1389	7	0
1	B	1332	1377	1374	3	1
1	C	1271	1314	1311	1	1
2	A	38	0	0	0	0
2	B	25	0	0	0	0
2	C	11	0	0	0	0
All	All	4025	4083	4074	10	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:LYS:HG2	1:A:250:TRP:CE2	2.40	0.57
1:A:206:LYS:HG2	1:A:250:TRP:CZ2	2.41	0.55
1:C:184:SER:N	1:C:185:PRO:CD	2.71	0.54
1:B:139:ILE:HD11	1:B:150:ALA:HB2	1.91	0.52
1:A:243:ARG:NH2	1:B:199:LEU:O	2.47	0.47
1:A:152:SER:HB3	1:A:158:VAL:HG23	1.98	0.45
1:B:161:PHE:CD2	1:B:178:ILE:HD12	2.54	0.42
1:A:234:TYR:CE2	1:A:235:ARG:HG3	2.54	0.42
1:A:285:ARG:O	1:A:289:GLU:HG3	2.21	0.41
1:A:285:ARG:HA	1:A:285:ARG:HD3	1.90	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:149:ARG:NH2	1:C:257:GLU:O[8_555]	2.08	0.12

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	169/171 (99%)	163 (96%)	6 (4%)	0	100	100
1	B	166/171 (97%)	160 (96%)	6 (4%)	0	100	100
1	C	152/171 (89%)	145 (95%)	7 (5%)	0	100	100
All	All	487/513 (95%)	468 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/146 (100%)	146 (100%)	0	100	100
1	B	145/146 (99%)	144 (99%)	1 (1%)	84	94
1	C	140/146 (96%)	139 (99%)	1 (1%)	84	94
All	All	431/438 (98%)	429 (100%)	2 (0%)	88	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	184	SER
1	C	209	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	171/171 (100%)	0.32	2 (1%)	79	80	26, 41, 72, 110	0
1	B	168/171 (98%)	0.45	10 (5%)	21	22	23, 41, 97, 132	0
1	C	158/171 (92%)	1.11	31 (19%)	1	1	30, 62, 98, 133	0
All	All	497/513 (96%)	0.61	43 (8%)	10	10	23, 46, 95, 133	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	135	MET	10.9
1	C	158	VAL	6.1
1	C	157	LYS	6.1
1	C	156	ASN	5.9
1	C	137	VAL	5.9
1	B	182	ALA	5.6
1	C	159	ARG	5.2
1	B	183	GLU	4.9
1	C	178	ILE	4.6
1	A	154	LYS	4.1
1	C	175	ILE	3.6
1	C	285	ARG	3.5
1	C	187	ALA	3.5
1	C	179	LEU	3.5
1	B	180	GLY	3.4
1	C	219	THR	3.3
1	C	176	PRO	3.2
1	C	177	THR	3.2
1	B	152	SER	3.2
1	C	161	PHE	3.1
1	C	138	ALA	3.0
1	C	225	ILE	2.9
1	C	218	ARG	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	216	VAL	2.8
1	C	300	ARG	2.8
1	C	295	ILE	2.7
1	C	184	SER	2.7
1	B	181	ALA	2.7
1	C	139	ILE	2.6
1	C	140	ILE	2.6
1	B	274	ARG	2.4
1	C	213	ILE	2.4
1	B	184	SER	2.3
1	B	234	TYR	2.2
1	B	285	ARG	2.2
1	C	150	ALA	2.2
1	A	302	ILE	2.2
1	C	261	LEU	2.1
1	C	149	ARG	2.1
1	C	186	GLY	2.1
1	B	159	ARG	2.1
1	C	299	LEU	2.0
1	C	200	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.