



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 08:50 am BST

PDB ID : 4CX2
Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex with 6-(5-(((3R,4R)-4-((6-AMINO-4-METHYLPYRIDIN-2-YL)METHYL) P YRROLIDIN-3-YL)OXY)PENTYL)-4-METHYLPYRIDIN-2-AMINE
Authors : Li, H.; Poulos, T.L.
Deposited on : 2014-04-03
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	 4% 82% 9% 9%
1	B	443	 2% 81% 9% 9%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7117 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	As	C	N	O	S			
1	A	405	3229	1	2053	569	590	16	0	1	0
1	B	403	3212	1	2042	566	587	16	0	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	conflict	UNP P29473
A	111	ALA	LEU	engineered mutation	UNP P29473
B	100	ARG	CYS	conflict	UNP P29473
B	111	ALA	LEU	engineered mutation	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 2-AMINO-6-(1,2-DIHYDROXY-PROPYL)-7,8-DIHYDRO-6H-PTERIDIN-4-ONE (three-letter code: H2B) (formula: C₉H₁₃N₅O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 6-[5-((3R,4R)-4-[(6-amino-4-methylpyridin-2-yl)methyl]pyrrolidin-3-yl)oxy]pentyl]-4-methylpyridin-2-amine (three-letter code: HW1) (formula: C₂₂H₃₃N₅O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
4	A	1	28	22	5	1	0	0
4	B	1	28	22	5	1	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	2	2	0	0
5	A	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	256	Total	O	0	0
			256	256		
7	B	227	Total	O	0	0
			227	227		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



- Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.88Å 106.57Å 156.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.13 – 2.04 48.38 – 2.04	Depositor EDS
% Data completeness (in resolution range)	95.2 (88.13-2.04) 95.3 (48.38-2.04)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 2.03Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.163 , 0.201 0.162 , 0.201	Depositor DCC
R_{free} test set	2972 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtrriage
Anisotropy	0.289	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7117	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CAS, ACT, HEM, HW1, H2B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.62	5/3312 (0.2%)	0.63	1/4509 (0.0%)
1	B	0.63	4/3291 (0.1%)	0.63	2/4480 (0.0%)
All	All	0.62	9/6603 (0.1%)	0.63	3/8989 (0.0%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	324	TRP	CD2-CE2	5.63	1.48	1.41
1	A	447	TRP	CD2-CE2	5.49	1.48	1.41
1	A	281	TRP	CD2-CE2	5.47	1.48	1.41
1	A	358	TRP	CD2-CE2	5.42	1.47	1.41
1	B	281	TRP	CD2-CE2	5.36	1.47	1.41
1	B	447	TRP	CD2-CE2	5.21	1.47	1.41
1	B	396	TRP	CD2-CE2	5.12	1.47	1.41
1	B	180	TRP	CD2-CE2	5.11	1.47	1.41
1	A	192	TRP	CD2-CE2	5.04	1.47	1.41

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	240	ARG	NE-CZ-NH2	-6.14	117.23	120.30
1	B	240	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	294	LEU	CA-CB-CG	5.07	126.96	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3229	0	3135	21	0
1	B	3212	0	3117	21	0
2	A	43	0	30	3	0
2	B	43	0	30	5	0
3	A	17	0	13	1	0
3	B	17	0	13	1	0
4	A	28	0	33	2	0
4	B	28	0	33	2	0
5	A	8	0	6	0	0
5	B	8	0	6	0	0
6	B	1	0	0	0	0
7	A	256	0	0	1	0
7	B	227	0	0	1	0
All	All	7117	0	6416	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:CAS:AS	1:A:384:CAS:SG	2.46	1.34
1:B:384:CAS:AS	1:B:384:CAS:SG	2.55	1.24
2:B:500:HEM:HBC2	2:B:500:HEM:HMC2	1.60	0.82
1:A:257:ARG:HG3	1:A:257:ARG:HH11	1.48	0.78
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.68	0.76
1:A:294:LEU:HD13	1:A:304:LEU:HD13	1.78	0.65
1:B:236:ARG:HD2	1:B:242:ASP:OD1	1.98	0.64
1:A:257:ARG:NH1	1:A:257:ARG:HG3	2.17	0.59
1:B:358:TRP:H	2:B:500:HEM:HAB	1.67	0.59
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.86	0.57
1:B:240:ARG:HD2	1:B:241:GLY:O	2.06	0.56
2:B:500:HEM:HBC2	2:B:500:HEM:CMC	2.31	0.56
1:A:384:CAS:CB	1:A:384:CAS:AS	3.14	0.55
1:B:240:ARG:HD3	1:B:298:PRO:HB3	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:LEU:HD11	1:A:156:GLU:HA	1.91	0.53
1:B:378:LEU:HB2	7:B:2179:HOH:O	2.07	0.53
1:A:378:LEU:HB2	7:A:2194:HOH:O	2.06	0.53
1:A:121:PRO:O	1:A:124:GLN:HB3	2.09	0.53
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.49	0.52
2:B:500:HEM:C1C	4:B:800:HW1:H34	2.45	0.52
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.51	0.51
1:B:449:TRP:HA	3:B:600:H2B:N1	2.25	0.51
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.42	0.50
1:B:455:SER:HB3	1:B:458:LEU:HD22	1.92	0.50
1:B:308:PRO:HD2	1:B:311:LEU:HD12	1.93	0.50
1:B:281:TRP:HB2	1:B:304:LEU:HD21	1.94	0.49
1:A:449:TRP:HA	3:A:600:H2B:N1	2.28	0.49
1:A:366:THR:O	1:A:370:CYS:HB2	2.14	0.47
1:B:366:THR:O	1:B:370:CYS:HB2	2.14	0.47
1:A:281:TRP:HB2	1:A:304:LEU:HD21	1.97	0.47
1:A:325:PHE:O	1:A:328:LEU:HB2	2.14	0.46
1:B:231:THR:O	1:B:353:ALA:HA	2.15	0.46
1:B:105:LEU:HD12	1:B:109:ARG:HD3	1.98	0.45
2:A:500:HEM:O1D	4:A:800:HW1:N01	2.50	0.44
1:A:452:PRO:HG2	1:A:459:THR:HG21	2.00	0.44
2:A:500:HEM:CBB	2:A:500:HEM:HHC	2.48	0.44
1:A:338:VAL:HG23	4:A:800:HW1:H31	2.00	0.44
1:B:167:LEU:HG	1:B:348:LEU:HD12	2.01	0.43
1:B:172:LEU:HD11	1:B:232:VAL:HG11	2.00	0.43
1:B:71:PRO:HG2	1:B:84:ASP:HB3	2.01	0.43
2:A:500:HEM:HBB2	2:A:500:HEM:HHC	2.01	0.42
1:A:423:HIS:HB2	1:B:392:THR:HB	2.02	0.41
1:A:392:THR:HB	1:B:423:HIS:HB2	2.02	0.41
1:B:178:GLN:HE22	1:B:181:ARG:HH11	1.67	0.41
2:B:500:HEM:NC	4:B:800:HW1:H34	2.36	0.40
1:A:472:SER:HA	1:A:473:PRO:C	2.42	0.40
1:A:481:PRO:HD2	1:A:482:TRP:CZ3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/443 (90%)	389 (97%)	12 (3%)	0	100	100
1	B	398/443 (90%)	387 (97%)	11 (3%)	0	100	100
All	All	799/886 (90%)	776 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/374 (92%)	330 (96%)	14 (4%)	30	23
1	B	342/374 (91%)	330 (96%)	12 (4%)	36	29
All	All	686/748 (92%)	660 (96%)	26 (4%)	33	26

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	72	ARG
1	A	99	ARG
1	A	100	ARG
1	A	102	LEU
1	A	123	GLU
1	A	125	LEU
1	A	154	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	398	ASP
1	A	458	LEU
1	A	468	ASN
1	B	99	ARG
1	B	100	ARG
1	B	102	LEU
1	B	105	LEU
1	B	223	ARG
1	B	225	ASN
1	B	240	ARG
1	B	330	LEU
1	B	378	LEU
1	B	389	THR
1	B	414	LEU
1	B	458	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	153	GLN
1	A	166	HIS
1	A	191	GLN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	178	GLN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CAS	B	384	1	5,8,9	1.25	0	1,9,11	0.57	0
1	CAS	A	384	1	5,8,9	1.07	0	1,9,11	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	B	384	1	-	0/0/7/9	-
1	CAS	A	384	1	-	0/0/7/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	384	CAS	1	0
1	A	384	CAS	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACT	A	861	-	1,3,3	2.17	1 (100%)	0,3,3	0.00	-
2	HEM	A	500	1	27,50,50	2.36	10 (37%)	17,82,82	3.24	10 (58%)
2	HEM	B	500	1	27,50,50	2.38	11 (40%)	17,82,82	3.18	9 (52%)
5	ACT	B	861	-	1,3,3	1.66	0	0,3,3	0.00	-
4	HW1	B	800	-	29,30,30	0.65	0	34,40,40	2.13	10 (29%)
5	ACT	B	860	-	1,3,3	1.63	0	0,3,3	0.00	-
5	ACT	A	860	-	1,3,3	1.65	0	0,3,3	0.00	-
3	H2B	B	600	-	13,18,18	1.40	2 (15%)	11,26,26	1.86	3 (27%)
3	H2B	A	600	-	13,18,18	1.28	1 (7%)	11,26,26	2.05	4 (36%)
4	HW1	A	800	-	29,30,30	0.78	0	34,40,40	1.79	9 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1	-	0/6/54/54	-
2	HEM	B	500	1	-	0/6/54/54	-
4	HW1	B	800	-	-	0/13/23/23	0/3/3/3
3	H2B	B	600	-	-	0/8/33/33	0/2/2/2
3	H2B	A	600	-	-	0/8/33/33	0/2/2/2
4	HW1	A	800	-	-	0/13/23/23	0/3/3/3

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	C3C-C2C	6.63	1.49	1.40
2	B	500	HEM	C3C-C2C	5.50	1.48	1.40
2	B	500	HEM	C4A-NA	4.99	1.46	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	C4A-NA	4.69	1.45	1.36
2	B	500	HEM	C3B-C2B	4.25	1.46	1.40
2	B	500	HEM	C1A-NA	4.09	1.44	1.36
3	B	600	H2B	C4A-N5	3.98	1.37	1.28
2	A	500	HEM	C1A-NA	3.82	1.44	1.36
2	B	500	HEM	C2A-C3A	3.68	1.48	1.37
2	A	500	HEM	C1C-C2C	3.43	1.50	1.42
2	A	500	HEM	C2A-C3A	3.42	1.47	1.37
2	B	500	HEM	C1C-C2C	3.19	1.49	1.42
2	A	500	HEM	C3B-C2B	3.15	1.44	1.40
3	A	600	H2B	C4A-N5	3.06	1.35	1.28
2	B	500	HEM	C3D-C2D	2.86	1.46	1.37
2	A	500	HEM	C3D-C2D	2.84	1.46	1.37
2	B	500	HEM	C1B-C2B	2.72	1.48	1.42
2	B	500	HEM	C4D-C3D	2.54	1.48	1.42
2	A	500	HEM	C4D-C3D	2.49	1.48	1.42
2	B	500	HEM	C4B-CHC	2.47	1.47	1.41
2	A	500	HEM	C1B-C2B	2.45	1.48	1.42
3	B	600	H2B	C8A-N1	2.31	1.37	1.32
2	B	500	HEM	C4A-CHB	2.18	1.47	1.41
5	A	861	ACT	CH3-C	2.17	1.51	1.48
2	A	500	HEM	C4B-CHC	2.00	1.46	1.41

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	HEM	C1D-C2D-C3D	-8.92	100.79	107.00
2	B	500	HEM	C1D-C2D-C3D	-8.75	100.91	107.00
4	B	800	HW1	C02-N01-C06	6.52	123.05	118.10
4	B	800	HW1	C22-N21-C26	5.26	122.08	118.10
2	B	500	HEM	CBA-CAA-C2A	-5.23	102.84	112.49
4	A	800	HW1	C02-N01-C06	4.95	121.85	118.10
2	A	500	HEM	C4A-C3A-C2A	-4.60	103.79	107.00
2	A	500	HEM	C3B-C4B-NB	4.54	115.08	109.21
2	B	500	HEM	C3B-C4B-NB	4.42	114.92	109.21
3	A	600	H2B	N1-C2-N3	-4.21	119.59	126.43
4	A	800	HW1	C22-N21-C26	4.16	121.25	118.10
3	A	600	H2B	C4-C4A-N5	3.90	122.34	118.06
3	B	600	H2B	C4-C4A-N5	3.84	122.28	118.06
2	A	500	HEM	CBA-CAA-C2A	-3.77	105.53	112.49
3	B	600	H2B	N1-C2-N3	-3.63	120.54	126.43
4	A	800	HW1	C5'-N1'-C2'	3.55	113.80	105.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	800	HW1	C5'-N1'-C2'	3.39	113.42	105.42
2	B	500	HEM	CBD-CAD-C3D	-3.34	106.33	112.48
4	B	800	HW1	C25-C26-N21	-3.26	119.44	122.90
4	B	800	HW1	C05-C06-N01	-3.07	119.64	122.90
2	A	500	HEM	CMB-C2B-C3B	3.07	130.42	124.68
4	B	800	HW1	C4'-C5'-N1'	-3.06	100.07	105.20
2	B	500	HEM	CMB-C2B-C3B	3.02	130.33	124.68
2	A	500	HEM	CBD-CAD-C3D	-2.97	107.01	112.48
4	A	800	HW1	C25-C26-N21	-2.96	119.76	122.90
4	A	800	HW1	C14-C26-N21	2.79	120.10	115.95
2	B	500	HEM	CMD-C2D-C3D	2.75	130.12	124.94
2	B	500	HEM	C4A-C3A-C2A	-2.73	105.09	107.00
4	A	800	HW1	C04-C05-C06	-2.64	118.59	120.32
4	A	800	HW1	C4'-C5'-N1'	-2.56	100.90	105.20
4	A	800	HW1	C05-C06-N01	-2.45	120.31	122.90
2	A	500	HEM	CAD-CBD-CGD	-2.41	108.63	112.67
2	B	500	HEM	CMC-C2C-C3C	2.22	128.83	124.68
3	A	600	H2B	N2-C2-N1	2.20	120.13	116.57
2	A	500	HEM	CMD-C2D-C3D	2.15	129.00	124.94
2	A	500	HEM	CAA-CBA-CGA	2.15	116.28	112.67
2	B	500	HEM	CAD-CBD-CGD	-2.12	109.11	112.67
2	A	500	HEM	CMA-C3A-C2A	2.09	128.89	124.94
4	B	800	HW1	C04-C05-C06	-2.09	118.95	120.32
4	B	800	HW1	C14-C26-N21	2.09	119.06	115.95
4	B	800	HW1	N22-C22-N21	2.08	119.78	116.49
4	B	800	HW1	C06-C08-C3'	-2.05	108.80	115.55
4	A	800	HW1	C06-C08-C3'	-2.04	108.83	115.55
3	A	600	H2B	N2-C2-N3	2.04	120.27	117.06
3	B	600	H2B	N2-C2-N1	2.03	119.87	116.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	3	0
2	B	500	HEM	5	0
4	B	800	HW1	2	0
3	B	600	H2B	1	0
3	A	600	H2B	1	0



Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	800	HW1	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/443 (91%)	0.07	18 (4%) 33 35	25, 36, 64, 91	0
1	B	402/443 (90%)	-0.11	10 (2%) 57 61	25, 40, 65, 98	0
All	All	806/886 (90%)	-0.02	28 (3%) 44 48	25, 38, 64, 98	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	GLN	6.3
1	A	109	ARG	5.2
1	A	259	GLN	5.1
1	A	67	GLY	5.1
1	B	261	GLY	5.0
1	A	160	ALA	4.5
1	A	69	LYS	4.0
1	A	123	GLU	3.5
1	A	121	PRO	3.5
1	B	142	ARG	3.0
1	B	260	ASP	2.9
1	A	239	GLY	2.8
1	B	146	GLN	2.8
1	A	449	TRP	2.7
1	B	109	ARG	2.7
1	A	450	ILE	2.5
1	B	144	GLY	2.4
1	A	152	LEU	2.4
1	A	142	ARG	2.4
1	A	124	GLN	2.3
1	A	238	PRO	2.3
1	A	146	GLN	2.3
1	B	448	ALA	2.2
1	A	257	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	323	GLU	2.1
1	B	239	GLY	2.1
1	A	126	LEU	2.1
1	A	153	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CAS	B	384	9/10	0.94	0.10	53,55,76,80	0
1	CAS	A	384	9/10	0.97	0.09	37,38,56,61	0

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	HW1	A	800	28/28	0.95	0.13	26,33,37,38	0
4	HW1	B	800	28/28	0.96	0.10	29,33,36,36	0
5	ACT	B	860	4/4	0.97	0.15	47,50,51,51	0
2	HEM	B	500	43/43	0.98	0.12	27,29,31,33	0
5	ACT	A	860	4/4	0.98	0.15	42,46,47,47	0
3	H2B	B	600	17/17	0.98	0.16	25,26,27,28	0
2	HEM	A	500	43/43	0.98	0.18	26,28,32,34	0
5	ACT	B	861	4/4	0.99	0.07	32,34,34,34	0
3	H2B	A	600	17/17	0.99	0.16	25,27,30,30	0
5	ACT	A	861	4/4	0.99	0.09	29,30,30,31	0
6	ZN	B	900	1/1	1.00	0.10	32,32,32,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEM B 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.