wwPDB X-ray Structure Validation Summary Report (i)

Jun 13, 2024 – 10:09 AM EDT

PDB ID	:	4D9E
Title	:	D-Cysteine desulfhydrase from Salmonella typhimurium complexed with L-cycloserine (LCS)
Authors	:	Bharath, S.R.; Shveta, B.; Rajesh, K.H.; Savithri, H.S.; Murthy, M.R.N.
Deposited on	:	2012-01-11
Resolution	:	2.47 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	80% 14% • • •
1	B	342	2% 79% 13% • 5%
1	C	342	3% 79% 12% • 6%
1	D	342	85% 10% • •

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 10065 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-Cysteine desulfhydrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	328	2435	1555	414	457	9	4	1	0
1	B	325	2414	1537	408	460	9	6	1	0
1	C	323	2419	1540	412	458	9	20	3	0
1	D	328	2434	1553	408	464	9	10	1	0

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	expression tag	UNP Q8ZNT7
A	-12	ARG	-	expression tag	UNP Q8ZNT7
A	-11	GLY	-	expression tag	UNP Q8ZNT7
A	-10	SER	-	expression tag	UNP Q8ZNT7
A	-9	HIS	-	expression tag	UNP Q8ZNT7
A	-8	HIS	-	expression tag	UNP Q8ZNT7
A	-7	HIS	-	expression tag	UNP Q8ZNT7
A	-6	HIS	-	expression tag	UNP Q8ZNT7
A	-5	HIS	-	expression tag	UNP Q8ZNT7
A	-4	HIS	-	expression tag	UNP Q8ZNT7
A	-3	GLY	-	expression tag	UNP Q8ZNT7
A	-2	MET	-	expression tag	UNP Q8ZNT7
A	-1	ALA	-	expression tag	UNP Q8ZNT7
A	0	SER	-	expression tag	UNP Q8ZNT7
B	-13	MET	-	expression tag	UNP Q8ZNT7
B	-12	ARG	-	expression tag	UNP Q8ZNT7
B	-11	GLY	-	expression tag	UNP Q8ZNT7
B	-10	SER	-	expression tag	UNP Q8ZNT7
B	-9	HIS	-	expression tag	UNP Q8ZNT7
B	-8	HIS	-	expression tag	UNP Q8ZNT7
B	-7	HIS	-	expression tag	UNP Q8ZNT7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	HIS	-	expression tag	UNP Q8ZNT7
B	-5	HIS	-	expression tag	UNP Q8ZNT7
B	-4	HIS	-	expression tag	UNP Q8ZNT7
B	-3	GLY	-	expression tag	UNP Q8ZNT7
B	-2	MET	-	expression tag	UNP Q8ZNT7
B	-1	ALA	-	expression tag	UNP Q8ZNT7
B	0	SER	-	expression tag	UNP Q8ZNT7
C	-13	MET	-	expression tag	UNP Q8ZNT7
C	-12	ARG	-	expression tag	UNP Q8ZNT7
C	-11	GLY	-	expression tag	UNP Q8ZNT7
C	-10	SER	-	expression tag	UNP Q8ZNT7
C	-9	HIS	-	expression tag	UNP Q8ZNT7
C	-8	HIS	-	expression tag	UNP Q8ZNT7
C	-7	HIS	-	expression tag	UNP Q8ZNT7
C	-6	HIS	-	expression tag	UNP Q8ZNT7
C	-5	HIS	-	expression tag	UNP Q8ZNT7
C	-4	HIS	-	expression tag	UNP Q8ZNT7
C	-3	GLY	-	expression tag	UNP Q8ZNT7
C	-2	MET	-	expression tag	UNP Q8ZNT7
C	-1	ALA	-	expression tag	UNP Q8ZNT7
C	0	SER	-	expression tag	UNP Q8ZNT7
D	-13	MET	-	expression tag	UNP Q8ZNT7
D	-12	ARG	-	expression tag	UNP Q8ZNT7
D	-11	GLY	-	expression tag	UNP Q8ZNT7
D	-10	SER	-	expression tag	UNP Q8ZNT7
D	-9	HIS	-	expression tag	UNP Q8ZNT7
D	-8	HIS	-	expression tag	UNP Q8ZNT7
D	-7	HIS	-	expression tag	UNP Q8ZNT7
D	-6	HIS	-	expression tag	UNP Q8ZNT7
D	-5	HIS	-	expression tag	UNP Q8ZNT7
D	-4	HIS	-	expression tag	UNP Q8ZNT7
D	-3	GLY	-	expression tag	UNP Q8ZNT7
D	-2	MET	-	expression tag	UNP Q8ZNT7
D	-1	ALA	-	expression tag	UNP Q8ZNT7
D	0	SER	-	expression tag	UNP Q8ZNT7

- Molecule 2 is BENZAMIDINE (three-letter code: BEN) (formula: $C_7H_8N_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N 9 7 2	0	0
2	B	1	Total C N 9 7 2	0	0
2	D	1	Total C N 9 7 2	0	0

- Molecule 3 is [5-hydroxy-6-methyl-4-({[(4E)-3-oxo-1,2-oxazolidin-4-ylidene]amino}methyl)pyridin-3-yl]methyl dihydrogen phosphate (three-letter code: LCS) (formula: $C_{11}H_{14}N_3O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	22	11	3	7	1	0	0
3	B	1	22	11	3	7	1	0	0
3	C	1	22	11	3	7	1	0	0
3	D	1	22	11	3	7	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	68	68	68	0	0
4	B	53	53	53	0	0
4	C	54	54	54	0	0
4	D	73	73	73	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: D-Cysteine desulfhydrase

Chain A:



• Molecule 1: D-Cysteine desulfhydrase

Chain B:



• Molecule 1: D-Cysteine desulfhydrase

Chain C:



• Molecule 1: D-Cysteine desulfhydrase

Chain D:


85%
10%
MET
ARG
GLY
SER
HIS
HIS
HIS
HIS
HIS
HIS
GLY
MET
ALA
SER
M1
P2
L3
L6
E13
R26
R33
I37
V42
L52
T72
H80
Q83
T105
E127
A166
C180
E181
E182
V183
V184
G185
A196
K228
I238
L242
A243
L244
I250
K273
L274
L275
L278
L282
R303
F304
N305
D306
H314
T315
G316
H325
V328

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	66.42Å 165.84Å 68.06Å 90.00° 119.03° 90.00°	Depositor
Resolution (Å)	58.08 – 2.47 58.08 – 2.47	Depositor EDS
% Data completeness (in resolution range)	90.1 (58.08-2.47) 90.1 (58.08-2.47)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.30 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.213 , 0.263 0.211 , 0.263	Depositor DCC
R_{free} test set	2121 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å^2)	25.6	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 37.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.29$	Xtriage
Estimated twinning fraction	0.023 for -h-l,k,h 0.023 for l,k,-h-l 0.044 for h,-k,-h-l 0.068 for -h-l,-k,l 0.046 for l,-k,h	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	10065	wwPDB-VP
Average B, all atoms (Å^2)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LCS, BEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.78	1/2482 (0.0%)	0.83	6/3385 (0.2%)
1	B	0.82	2/2460 (0.1%)	0.82	4/3354 (0.1%)
1	C	0.88	4/2474 (0.2%)	0.82	4/3371 (0.1%)
1	D	0.78	2/2484 (0.1%)	0.81	0/3389
All	All	0.82	9/9900 (0.1%)	0.82	14/13499 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	129	CYS	CB-SG	-8.44	1.67	1.82
1	B	180	CYS	CB-SG	-6.39	1.71	1.82
1	B	90	LYS	CB-CG	-5.94	1.36	1.52
1	C	180	CYS	CB-SG	-5.92	1.72	1.81
1	D	180	CYS	CB-SG	-5.77	1.72	1.81

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	184	VAL	N-CA-C	-7.78	89.99	111.00
1	A	184	VAL	N-CA-CB	-6.97	96.16	111.50
1	A	183	VAL	CB-CA-C	6.38	123.51	111.40
1	B	275	LEU	CD1-CG-CD2	-5.89	92.83	110.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	270	GLU	OE1-CD-OE2	-5.78	116.37	123.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	7	THR	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2435	0	2445	39	0
1	B	2414	0	2403	36	0
1	C	2419	0	2434	40	0
1	D	2434	0	2436	29	0
2	A	9	0	7	0	0
2	B	9	0	7	0	0
2	D	9	0	7	0	0
3	A	22	0	11	1	0
3	B	22	0	11	2	0
3	C	22	0	11	5	0
3	D	22	0	11	1	0
4	A	68	0	0	0	0
4	B	53	0	0	0	0
4	C	54	0	0	1	0
4	D	73	0	0	0	0
All	All	10065	0	9783	141	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:182:GLU:O	1:D:183:VAL:HG23	1.49	1.11

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:184:VAL:CB	1:B:185:GLY:HA2	1.90	0.99
1:A:183:VAL:HG12	1:A:184:VAL:HG12	1.42	0.98
1:A:182:GLU:O	1:A:183:VAL:HG23	1.74	0.87
1:D:183:VAL:O	1:D:183:VAL:HG12	1.73	0.87

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/342 (95%)	314 (96%)	9 (3%)	3 (1%)	17	29
1	B	323/342 (94%)	314 (97%)	7 (2%)	2 (1%)	25	40
1	C	324/342 (95%)	307 (95%)	13 (4%)	4 (1%)	13	22
1	D	327/342 (96%)	312 (95%)	10 (3%)	5 (2%)	10	16
All	All	1300/1368 (95%)	1247 (96%)	39 (3%)	14 (1%)	14	23

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	183	VAL
1	B	183	VAL
1	B	184	VAL
1	C	8	ARG
1	C	181	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/267 (94%)	240 (96%)	10 (4%)	31	53
1	B	248/267 (93%)	233 (94%)	15 (6%)	19	34
1	C	251/267 (94%)	236 (94%)	15 (6%)	19	34
1	D	251/267 (94%)	245 (98%)	6 (2%)	49	72
All	All	1000/1068 (94%)	954 (95%)	46 (5%)	27	47

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	64	ARG
1	C	183	VAL
1	C	65	GLU
1	C	142	LEU
1	C	303	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	83	GLN
1	C	314	HIS
1	D	314	HIS
1	C	178	GLN
1	C	325	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	BEN	D	401	-	9,9,9	1.24	0	7,11,11	1.99	2 (28%)
2	BEN	A	401	-	9,9,9	1.46	1 (11%)	7,11,11	0.93	0
3	LCS	D	402	-	20,23,23	3.78	13 (65%)	23,33,33	1.90	7 (30%)
2	BEN	B	401	-	9,9,9	1.08	1 (11%)	7,11,11	1.29	1 (14%)
3	LCS	B	402	-	20,23,23	4.02	11 (55%)	23,33,33	2.90	11 (47%)
3	LCS	C	401	-	20,23,23	4.42	9 (45%)	23,33,33	2.57	14 (60%)
3	LCS	A	402	-	20,23,23	4.02	13 (65%)	23,33,33	2.87	9 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEN	D	401	-	-	0/4/4/4	0/1/1/1
2	BEN	A	401	-	-	0/4/4/4	0/1/1/1
3	LCS	D	402	-	-	8/10/21/21	0/2/2/2
2	BEN	B	401	-	-	0/4/4/4	0/1/1/1
3	LCS	B	402	-	-	5/10/21/21	0/2/2/2
3	LCS	C	401	-	-	6/10/21/21	0/2/2/2
3	LCS	A	402	-	-	4/10/21/21	0/2/2/2

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	401	LCS	CA-N	13.30	1.46	1.28
3	B	402	LCS	CA-N	12.87	1.46	1.28
3	A	402	LCS	CA-N	11.61	1.44	1.28
3	D	402	LCS	CA-N	11.01	1.43	1.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	401	LCS	C3-C2	9.33	1.50	1.41

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	402	LCS	C4A-C4-C5	-9.00	110.00	119.76
3	A	402	LCS	C3-C4-C5	-8.45	111.07	118.73
3	A	402	LCS	C4A-C4-C5	6.68	127.00	119.76
3	C	401	LCS	C6-C5-C4	6.25	122.79	118.06
3	C	401	LCS	O3P-P-O4P	4.68	118.87	106.67

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	LCS	CB-CA-N-C4A
3	B	402	LCS	C3-C4-C4A-N
3	B	402	LCS	C5-C4-C4A-N
3	B	402	LCS	CB-CA-N-C4A
3	C	401	LCS	C5A-O4P-P-O1P

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	402	LCS	1	0
3	B	402	LCS	2	0
3	C	401	LCS	5	0
3	A	402	LCS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand LCS D 402



Ligand LCS B 402



Ligand LCS C 401



Ligand LCS A 402



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	328/342 (95%)	-0.24	0	100	100	11, 20, 40, 47	3 (0%)
1	B	325/342 (95%)	-0.09	8 (2%)	57	59	11, 20, 37, 47	2 (0%)
1	C	323/342 (94%)	-0.08	9 (2%)	53	55	11, 20, 38, 45	5 (1%)
1	D	328/342 (95%)	-0.22	1 (0%)	94	94	11, 20, 40, 47	2 (0%)
All	All	1304/1368 (95%)	-0.15	18 (1%)	75	77	11, 20, 40, 47	12 (0%)

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	243	ALA	4.4
1	D	184	VAL	3.2
1	B	183	VAL	2.9
1	B	305	ASN	2.9
1	C	240	GLY	2.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	BEN	A	401	9/9	0.92	0.17	10,14,18,19	0
3	LCS	B	402	22/22	0.92	0.16	19,21,21,24	6
3	LCS	C	401	22/22	0.93	0.18	18,21,22,23	6
3	LCS	A	402	22/22	0.94	0.15	19,21,22,23	0
3	LCS	D	402	22/22	0.94	0.16	18,21,21,23	6
2	BEN	B	401	9/9	0.95	0.19	19,20,21,21	0
2	BEN	D	401	9/9	0.97	0.13	10,11,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LCS B 402:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around LCS C 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LCS A 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around LCS D 402: 2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)	
	

6.5 Other polymers (i)

There are no such residues in this entry.