



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 12:51 pm BST

PDB ID : 6DXB  
Title : Crystal structure of chalcone synthase from Arabidopsis thaliana  
Authors : Liou, G.; Chiang, Y.C.; Wang, Y.; Weng, J.K.  
Deposited on : 2018-06-28  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	 96%
1	B	395	 96%
1	C	395	 95%
1	D	395	 95%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 25857 atoms, of which 12125 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chalcone synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	389	6058	1910	3038	521	571	18	0	4	0
1	C	388	6044	1907	3030	519	570	18	0	4	0
1	B	389	6043	1906	3031	519	569	18	0	3	0
1	D	388	6032	1903	3026	518	567	18	0	3	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	432	Total	O	0	0
			432	432		
2	C	429	Total	O	0	0
			429	429		
2	B	403	Total	O	0	0
			403	403		
2	D	416	Total	O	0	0
			416	416		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chalcone synthase



- Molecule 1: Chalcone synthase



- Molecule 1: Chalcone synthase



- Molecule 1: Chalcone synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.64Å 137.56Å 108.56Å 90.00° 95.59° 90.00°	Depositor
Resolution (Å)	38.68 – 1.55 38.68 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.5 (38.68-1.55) 98.5 (38.68-1.55)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.59 (at 1.55Å)	Xtrriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.141 , 0.163 0.142 , 0.164	Depositor DCC
$R_{free}$ test set	1963 reflections (0.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.2	Xtrriage
Anisotropy	0.217	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 44.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	25857	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/3072	0.63	1/4155 (0.0%)
1	B	0.45	0/3064	0.63	1/4144 (0.0%)
1	C	0.42	0/3066	0.63	3/4147 (0.1%)
1	D	0.43	0/3058	0.64	2/4136 (0.0%)
All	All	0.43	0/12260	0.63	7/16582 (0.0%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	142	MET	CG-SD-CE	-7.52	88.16	100.20
1	B	142	MET	CG-SD-CE	-6.89	89.18	100.20
1	C	142	MET	CG-SD-CE	-6.15	90.36	100.20
1	A	142	MET	CG-SD-CE	-6.09	90.45	100.20
1	C	73	ARG	NE-CZ-NH1	5.65	123.13	120.30
1	C	73	ARG	NE-CZ-NH2	-5.57	117.52	120.30
1	D	161	ARG	NE-CZ-NH2	-5.40	117.60	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3020	3038	3034	6	0
1	B	3012	3031	3029	9	0
1	C	3014	3030	3027	6	0
1	D	3006	3026	3024	4	0
2	A	432	0	0	4	0
2	B	403	0	0	6	0
2	C	429	0	0	3	0
2	D	416	0	0	1	0
All	All	13732	12125	12114	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:211:LEU:HG	2:B:485:HOH:O	1.99	0.62
1:A:63:ARG:NH2	2:A:403:HOH:O	2.37	0.53
1:B:63:ARG:NH2	2:B:402:HOH:O	2.41	0.53
1:C:276:ASP:N	2:C:405:HOH:O	2.42	0.51
1:A:393:VAL:HG12	2:A:416:HOH:O	2.11	0.51
1:D:62:LYS:HD2	2:D:573:HOH:O	2.11	0.50
1:B:271[B]:PHE:CZ	1:B:273:LEU:HD21	2.49	0.46
1:C:391:HIS:HD2	1:C:392:SER:O	1.99	0.45
1:B:63:ARG:NE	2:B:402:HOH:O	2.51	0.44
1:C:275:LYS:HA	2:C:405:HOH:O	2.18	0.43
1:C:294:LYS:N	1:C:295:PRO:CD	2.82	0.43
1:A:48:ILE:HD13	1:A:79:GLU:HG3	2.01	0.43
1:D:48:ILE:HD13	1:D:79:GLU:HG3	2.00	0.43
1:A:294:LYS:N	1:A:295:PRO:CD	2.82	0.42
1:D:91:TYR:CD2	1:D:92:MET:HG2	2.54	0.42
1:B:358:LYS:HE2	2:B:478:HOH:O	2.19	0.42
1:B:63:ARG:CZ	2:B:402:HOH:O	2.68	0.41
1:C:91:TYR:CD2	1:C:92:MET:HG2	2.55	0.41
1:B:258:GLY:O	1:B:274:LEU:HB2	2.20	0.41
1:C:391:HIS:HE1	2:C:630:HOH:O	2.03	0.41
1:A:240:LYS:NZ	2:A:416:HOH:O	2.50	0.41
1:A:393:VAL:CG1	2:A:416:HOH:O	2.67	0.41
1:D:92:MET:SD	1:D:267:VAL:HG21	2.61	0.40
1:B:271[B]:PHE:HB3	2:B:483:HOH:O	2.22	0.40
1:B:193:VAL:O	1:B:226:ALA:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/395 (99%)	383 (98%)	7 (2%)	0	100	100
1	B	389/395 (98%)	380 (98%)	9 (2%)	0	100	100
1	C	389/395 (98%)	382 (98%)	7 (2%)	0	100	100
1	D	388/395 (98%)	379 (98%)	9 (2%)	0	100	100
All	All	1556/1580 (98%)	1524 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/327 (100%)	327 (100%)	1 (0%)	92	84
1	B	327/327 (100%)	326 (100%)	1 (0%)	92	84
1	C	327/327 (100%)	326 (100%)	1 (0%)	92	84
1	D	326/327 (100%)	323 (99%)	3 (1%)	78	60
All	All	1308/1308 (100%)	1302 (100%)	6 (0%)	88	77

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	356	ARG
1	C	356	ARG
1	B	356	ARG
1	D	87	HIS
1	D	157	PRO
1	D	356	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	391	HIS
1	D	391	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSD	D	169	1	3,7,8	0.73	0	1,8,10	1.44	0
1	CSD	A	169	1	3,7,8	0.94	0	1,8,10	1.43	0
1	CSD	C	169	1	3,7,8	0.87	0	1,8,10	1.03	0
1	CSD	B	169	1	3,7,8	0.74	0	1,8,10	2.26	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	D	169	1	-	1/2/6/8	-
1	CSD	A	169	1	-	1/2/6/8	-
1	CSD	C	169	1	-	1/2/6/8	-
1	CSD	B	169	1	-	1/2/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	169	CSD	OD1-SG-CB	-2.26	101.23	105.54

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	169	CSD	CA-CB-SG-OD1
1	A	169	CSD	CA-CB-SG-OD1
1	C	169	CSD	CA-CB-SG-OD1
1	B	169	CSD	CA-CB-SG-OD1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/395 (98%)	-0.38	3 (0%) 86 88	7, 13, 32, 50	0
1	B	388/395 (98%)	-0.30	9 (2%) 60 66	8, 13, 36, 76	0
1	C	387/395 (97%)	-0.32	1 (0%) 94 95	7, 13, 33, 52	0
1	D	387/395 (97%)	-0.39	1 (0%) 94 95	8, 12, 33, 59	0
All	All	1550/1580 (98%)	-0.35	14 (0%) 84 86	7, 13, 34, 76	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	211	LEU	13.3
1	B	210	HIS	5.1
1	B	209	THR	4.6
1	B	212	ASP	4.4
1	A	271[A]	PHE	3.6
1	A	274	LEU	3.3
1	B	208	ASP	3.0
1	B	271[A]	PHE	2.9
1	B	213	SER	2.8
1	A	7	SER	2.7
1	B	7	SER	2.4
1	C	55	THR	2.3
1	D	271[A]	PHE	2.2
1	B	274	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSD	B	169	8/9	0.96	0.09	11,14,23,28	0
1	CSD	A	169	8/9	0.97	0.09	11,13,26,31	0
1	CSD	C	169	8/9	0.97	0.10	10,14,32,39	0
1	CSD	D	169	8/9	0.97	0.08	9,12,29,35	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.