



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2023 – 10:58 PM EDT

PDB ID : 8F0L  
Title : Crystal Structure of the Human T cell Receptor CD3(EPSILON) N-Terminal Peptide Complexed with ADI-26906 FAB  
Authors : Battles, M.B.; Welin, M.  
Deposited on : 2022-11-03  
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	 88% 6% • 5%
1	H	223	 89% 5% • 5%
2	B	219	 91% 9%
2	L	219	 95% • •
3	P	13	 46% 8% 46%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	Q	13	 38% 15% 46%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	301	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14372 atoms, of which 6762 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADI-26906 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	H	212	Total 3239	C 1039	H 1603	N 274	O 314	S 9	0	3	0
1	A	212	Total 3233	C 1038	H 1599	N 272	O 315	S 9	0	3	0


- Molecule 2 is a protein called ADI-26906 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	L	216	Total 3466	C 1086	H 1729	N 297	O 347	S 7	0	14	0
2	B	218	Total 3472	C 1090	H 1731	N 297	O 348	S 6	0	11	0

- Molecule 3 is a protein called T-cell surface glycoprotein CD3 epsilon chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
3	P	7	Total 94	C 30	H 40	N 8	O 15	S 1	0	0	0
3	Q	7	Total 94	C 30	H 40	N 8	O 15	S 1	0	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	H	1	Total	C	H	O	0	0
			9	3	3	3		
4	H	1	Total	C	H	O	0	0
			9	3	3	3		
4	H	1	Total	C	H	O	0	0
			9	3	3	3		
4	L	1	Total	C	H	O	0	0
			8	3	2	3		
4	L	1	Total	C	H	O	0	0
			9	3	3	3		
4	A	1	Total	C	H	O	0	0
			9	3	3	3		
4	B	1	Total	C	H	O	0	0
			9	3	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	227	Total	O	0	0
			227	227		
5	L	191	Total	O	0	0
			191	191		
5	A	152	Total	O	0	0
			152	152		
5	B	136	Total	O	0	0
			136	136		
5	P	4	Total	O	0	0
			4	4		

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	Q	2	Total	O	0	0
			2	2		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ADI-26906 Fab Heavy Chain

Chain H:  89% 5% • 5%



- Molecule 1: ADI-26906 Fab Heavy Chain

Chain A:  88% 6% • 5%



- Molecule 2: ADI-26906 Fab Light Chain

Chain L:  95% • •




- Molecule 2: ADI-26906 Fab Light Chain

Chain B:  91% 9%



- Molecule 3: T-cell surface glycoprotein CD3 epsilon chain

Chain P:  46% 8% 46%



- Molecule 3: T-cell surface glycoprotein CD3 epsilon chain

Chain Q:  38% 15% 46%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.63Å 53.08Å 113.93Å 90.00° 99.82° 90.00°	Depositor
Resolution (Å)	48.60 – 1.81 48.59 – 1.81	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.60-1.81) 99.9 (48.59-1.81)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 1.81Å)	Xtrriage
Refinement program	PHENIX v1.19.2	Depositor
R, $R_{free}$	0.166 , 0.195 0.160 , 0.193	Depositor DCC
$R_{free}$ test set	3991 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.7	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	14372	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.91	6/1676 (0.4%)	0.78	0/2285
1	H	0.96	3/1678 (0.2%)	0.91	2/2287 (0.1%)
2	B	0.82	0/1808	0.79	1/2448 (0.0%)
2	L	0.86	3/1818 (0.2%)	0.82	0/2462
3	P	0.87	0/46	0.70	0/61
3	Q	0.86	0/46	0.74	0/61
All	All	0.89	12/7072 (0.2%)	0.82	3/9604 (0.0%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	152	GLU	CD-OE1	7.22	1.33	1.25
1	A	85	GLU	CB-CG	-7.02	1.38	1.52
2	L	92	TYR	CD2-CE2	6.51	1.49	1.39
2	L	36	TYR	CD1-CE1	-6.03	1.30	1.39
1	A	91	TYR	CD2-CE2	5.74	1.48	1.39
1	H	33	TYR	CE2-CZ	-5.46	1.31	1.38
1	A	85	GLU	CD-OE1	-5.39	1.19	1.25
2	L	92	TYR	CD1-CE1	5.25	1.47	1.39
1	H	84	SER	CB-OG	-5.19	1.35	1.42
1	A	144[A]	CYS	CB-SG	-5.08	1.73	1.81
1	A	144[B]	CYS	CB-SG	-5.08	1.73	1.81
1	H	192	SER	CB-OG	5.06	1.48	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	182	LEU	CA-CB-CG	6.39	129.99	115.30
1	H	101	ASP	CB-CG-OD1	5.27	123.04	118.30
2	B	170	ASP	CB-CG-OD1	5.03	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1634	1599	1598	8	0
1	H	1636	1603	1603	4	0
2	B	1741	1731	1734	12	0
2	L	1737	1729	1725	6	0
3	P	54	40	40	0	0
3	Q	54	40	40	1	0
4	A	6	3	7	6	0
4	B	6	3	7	1	0
4	H	18	9	23	0	0
4	L	12	5	14	1	0
5	A	152	0	0	5	0
5	B	136	0	0	0	0
5	H	227	0	0	0	0
5	L	191	0	0	5	0
5	P	4	0	0	0	0
5	Q	2	0	0	0	0
All	All	7610	6762	6791	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:142[B]:ARG:NH1	5:L:401:HOH:O	1.86	1.07
2:L:142[A]:ARG:NH1	5:L:401:HOH:O	2.05	0.90
1:A:1:PCA:N	5:A:401:HOH:O	2.08	0.85
1:H:147:LYS:HE2	5:L:422:HOH:O	1.99	0.63
2:L:138[A]:ASN:ND2	2:L:170:ASP:OD2	2.32	0.63
1:H:52(A):LEU:O	1:H:71[B]:ARG:NH1	2.32	0.62
4:L:301:GOL:H31	5:L:451:HOH:O	1.99	0.62

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:33:LEU:HD22	2:B:71:PHE:CG	2.35	0.61
2:B:32:TYR:OH	3:Q:6:GLU:OE1	2.17	0.60
2:B:183:LYS:O	2:B:187:GLU:HG2	2.01	0.60
4:A:301:GOL:H11	5:A:407:HOH:O	2.00	0.60
2:B:33:LEU:HG	2:B:34:ALA:N	2.21	0.55
1:A:182:LEU:HA	4:A:301:GOL:H12	1.89	0.55
2:B:105:GLU:OE1	2:B:173:TYR:OH	2.11	0.55
2:B:81:GLU:HG2	4:B:301:GOL:H11	1.90	0.54
4:A:301:GOL:H31	2:B:176[A]:SER:OG	2.11	0.51
2:L:13:VAL:HG13	2:L:17:GLU:HB2	1.93	0.51
4:A:301:GOL:C1	5:A:407:HOH:O	2.58	0.50
1:A:173:VAL:HG22	4:A:301:GOL:O1	2.12	0.50
2:L:201:LEU:HD13	2:L:205:VAL:HG12	1.94	0.49
2:B:33:LEU:HD22	2:B:71:PHE:CB	2.42	0.49
2:B:13:VAL:HG13	2:B:17:GLU:HB2	1.98	0.45
1:H:66:ARG:HD2	1:H:82(A):SER:O	2.18	0.43
1:A:183:SER:OG	4:A:301:GOL:H32	2.19	0.43
2:L:138[A]:ASN:ND2	5:L:404:HOH:O	2.41	0.43
1:A:217:PRO:C	5:A:403:HOH:O	2.58	0.42
2:B:145:LYS:HB3	2:B:197:THR:HB	2.02	0.42
1:H:36:TRP:CE2	1:H:80:MET:HB2	2.54	0.42
2:B:30(A):ASN:OD1	2:B:30(C):ARG:HG2	2.20	0.41
1:A:3[A]:GLN:NE2	5:A:406:HOH:O	2.44	0.41
1:A:50:TRP:C	1:A:50:TRP:CD1	2.94	0.41
2:B:37:GLN:HB2	2:B:47:LEU:HD11	2.02	0.41
1:A:204:HIS:HB3	1:A:209:THR:HB	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/223 (95%)	208 (99%)	3 (1%)	0	100	100
1	H	211/223 (95%)	209 (99%)	2 (1%)	0	100	100
2	B	227/219 (104%)	219 (96%)	8 (4%)	0	100	100
2	L	229/219 (105%)	223 (97%)	6 (3%)	0	100	100
3	P	5/13 (38%)	5 (100%)	0	0	100	100
3	Q	5/13 (38%)	5 (100%)	0	0	100	100
All	All	888/910 (98%)	869 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/188 (96%)	177 (98%)	4 (2%)	52	39
1	H	181/188 (96%)	178 (98%)	3 (2%)	60	50
2	B	201/191 (105%)	201 (100%)	0	100	100
2	L	204/191 (107%)	204 (100%)	0	100	100
3	P	5/9 (56%)	5 (100%)	0	100	100
3	Q	5/9 (56%)	5 (100%)	0	100	100
All	All	777/776 (100%)	770 (99%)	7 (1%)	81	74

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	71[A]	ARG
1	H	71[B]	ARG
1	H	182	LEU
1	A	50	TRP
1	A	71	ARG
1	A	83	ARG
1	A	153	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PCA	Q	1	3	7,8,9	2.34	2 (28%)	9,10,12	1.73	3 (33%)
1	PCA	A	1	1	7,8,9	2.07	2 (28%)	9,10,12	2.14	4 (44%)
3	PCA	P	1	3	7,8,9	2.98	2 (28%)	9,10,12	2.39	6 (66%)
1	PCA	H	1	1	7,8,9	2.05	2 (28%)	9,10,12	2.09	6 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PCA	Q	1	3	-	0/0/11/13	0/1/1/1
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1
3	PCA	P	1	3	-	0/0/11/13	0/1/1/1
1	PCA	H	1	1	-	0/0/11/13	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	1	PCA	CA-N	5.60	1.53	1.46
3	P	1	PCA	CD-N	4.91	1.47	1.34
1	A	1	PCA	CD-N	4.69	1.47	1.34

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	Q	1	PCA	CD-N	4.34	1.46	1.34
1	H	1	PCA	CD-N	4.27	1.45	1.34
3	Q	1	PCA	CA-N	4.05	1.51	1.46
1	A	1	PCA	CA-N	2.60	1.49	1.46
1	H	1	PCA	CA-N	2.36	1.49	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	1	PCA	CA-N-CD	-3.74	100.77	113.58
1	A	1	PCA	CA-N-CD	-3.50	101.60	113.58
3	Q	1	PCA	CA-N-CD	-3.26	102.43	113.58
1	A	1	PCA	CB-CA-N	3.16	112.36	103.30
3	P	1	PCA	OE-CD-CG	-3.07	121.40	126.76
1	H	1	PCA	CA-N-CD	-3.04	103.17	113.58
3	P	1	PCA	CG-CD-N	3.02	116.21	108.39
1	H	1	PCA	CB-CA-N	2.94	111.74	103.30
1	A	1	PCA	OE-CD-CG	-2.64	122.15	126.76
3	P	1	PCA	O-C-CA	-2.50	118.23	124.78
3	Q	1	PCA	CB-CA-N	2.45	110.32	103.30
3	P	1	PCA	CB-CA-N	2.40	110.20	103.30
1	A	1	PCA	CG-CD-N	2.40	114.61	108.39
1	H	1	PCA	OE-CD-CG	-2.30	122.75	126.76
1	H	1	PCA	CG-CD-N	2.29	114.32	108.39
3	P	1	PCA	CB-CG-CD	-2.17	100.90	104.40
1	H	1	PCA	O-C-CA	-2.05	119.39	124.78
1	H	1	PCA	CB-CA-C	-2.05	109.89	112.70
3	Q	1	PCA	CG-CD-N	2.03	113.65	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1	PCA	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	L	301	-	5,5,5	2.50	2 (40%)	5,5,5	0.61	0
4	GOL	H	303	-	5,5,5	1.48	1 (20%)	5,5,5	0.96	0
4	GOL	A	301	-	5,5,5	2.20	2 (40%)	5,5,5	1.51	1 (20%)
4	GOL	L	302	-	5,5,5	1.25	0	5,5,5	0.90	0
4	GOL	H	301	-	5,5,5	1.68	2 (40%)	5,5,5	0.75	0
4	GOL	B	301	-	5,5,5	1.34	1 (20%)	5,5,5	0.93	0
4	GOL	H	302	-	5,5,5	1.24	0	5,5,5	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	L	301	-	-	0/4/4/4	-
4	GOL	H	303	-	-	2/4/4/4	-
4	GOL	A	301	-	-	2/4/4/4	-
4	GOL	L	302	-	-	4/4/4/4	-
4	GOL	H	301	-	-	0/4/4/4	-
4	GOL	B	301	-	-	2/4/4/4	-
4	GOL	H	302	-	-	2/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	L	301	GOL	O3-C3	-4.60	1.23	1.42
4	A	301	GOL	O2-C2	3.51	1.53	1.43
4	H	301	GOL	C3-C2	2.87	1.63	1.51
4	L	301	GOL	O2-C2	-2.63	1.35	1.43

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	301	GOL	O1-C1	2.38	1.52	1.42
4	A	301	GOL	O3-C3	2.31	1.52	1.42
4	H	303	GOL	C1-C2	2.24	1.61	1.51
4	B	301	GOL	O2-C2	-2.21	1.36	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	301	GOL	C3-C2-C1	-2.78	100.89	111.70

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	302	GOL	C1-C2-C3-O3
4	H	303	GOL	O1-C1-C2-C3
4	L	302	GOL	O1-C1-C2-C3
4	A	301	GOL	O1-C1-C2-O2
4	A	301	GOL	O1-C1-C2-C3
4	H	302	GOL	O2-C2-C3-O3
4	L	302	GOL	C1-C2-C3-O3
4	B	301	GOL	O1-C1-C2-C3
4	L	302	GOL	O1-C1-C2-O2
4	B	301	GOL	O1-C1-C2-O2
4	H	303	GOL	O1-C1-C2-O2
4	L	302	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	301	GOL	1	0
4	A	301	GOL	6	0
4	B	301	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	211/223 (94%)	-0.37	1 (0%) 91 89	20, 33, 59, 72	0
1	H	211/223 (94%)	-0.51	0 100 100	17, 24, 41, 50	0
2	B	218/219 (99%)	-0.37	1 (0%) 91 89	23, 35, 60, 79	0
2	L	216/219 (98%)	-0.49	0 100 100	18, 29, 44, 56	0
3	P	6/13 (46%)	-0.24	0 100 100	28, 33, 43, 59	0
3	Q	6/13 (46%)	0.20	0 100 100	36, 46, 56, 75	0
All	All	868/910 (95%)	-0.43	2 (0%) 95 93	17, 30, 53, 79	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	205	LYS	2.1
2	B	126	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PCA	A	1	8/9	0.85	0.19	51,68,80,80	0
1	PCA	H	1	8/9	0.97	0.07	22,29,34,34	0
3	PCA	Q	1	8/9	0.97	0.07	28,32,39,39	0
3	PCA	P	1	8/9	0.99	0.06	20,24,29,29	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	301	6/6	0.81	0.23	33,40,48,48	0
4	GOL	H	303	6/6	0.82	0.20	34,38,55,55	0
4	GOL	B	301	6/6	0.82	0.16	40,53,64,67	0
4	GOL	L	301	6/6	0.89	0.15	30,39,43,47	0
4	GOL	H	302	6/6	0.92	0.19	33,35,46,46	0
4	GOL	L	302	6/6	0.93	0.25	31,37,50,52	0
4	GOL	H	301	6/6	0.97	0.05	22,24,29,29	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.