



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 12:09 PM BST

PDB ID : 3G70  
Title : Design and Preparation of Potent, Non-Peptidic, Bioavailable Renin Inhibitors  
Authors : Bezencon, O.; Bur, D.; Prade, L.; Weller, T.; Boss, C.; Fischli, W.  
Deposited on : 2009-02-09  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	 3% 82% 15% ..
1	B	341	 3% 78% 19% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DMS	B	343	-	-	X	-

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5706 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Renin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	334	Total 2593	C 1656	N 422	O 501	S 14	0	2	0
1	B	337	Total 2617	C 1673	N 425	O 505	S 14	0	2	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	341	HIS	-	expression tag	UNP P00797
B	341	HIS	-	expression tag	UNP P00797

- Molecule 2 is (1R,5S)-7-{4-[3-(2-chloro-3,6-difluorophenoxy)propyl]phenyl}-N-cyclopropyl-N-(2,3-dichlorobenzyl)-3,9-diazabicyclo[3.3.1]non-6-ene-6-carboxamide (three-letter code: A5T) (formula: C<sub>33</sub>H<sub>32</sub>Cl<sub>3</sub>F<sub>2</sub>N<sub>3</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	Cl	F	N	O	0	0
			43	33	3	2	3	2		
2	B	1	Total	C	Cl	F	N	O	0	0
			43	33	3	2	3	2		

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	175	175	175	0	0
5	B	199	199	199	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Renin



- Molecule 1: Renin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.73Å 89.46Å 118.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.98 – 2.00 36.98 – 2.00	Depositor EDS
% Data completeness (in resolution range)	84.4 (36.98-2.00) 84.4 (36.98-2.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.52 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.196 , 0.263 0.197 , 0.263	Depositor DCC
$R_{free}$ test set	2042 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.9	Xtrriage
Anisotropy	0.119	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 52.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5706	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, NAG, A5T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/2654	0.84	7/3598 (0.2%)
1	B	0.62	0/2678	0.85	8/3631 (0.2%)
All	All	0.61	0/5332	0.85	15/7229 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	165	ASP	CB-CG-OD2	6.42	124.08	118.30
1	B	226	ASP	CB-CG-OD2	6.41	124.07	118.30
1	B	38	ASP	CB-CG-OD2	6.37	124.04	118.30
1	A	145	ASP	CB-CG-OD2	6.05	123.75	118.30
1	B	283	ASP	CB-CG-OD2	5.88	123.60	118.30
1	B	219	ASP	CB-CG-OD2	5.70	123.43	118.30
1	A	283	ASP	CB-CG-OD2	5.68	123.41	118.30
1	B	254	ASP	CB-CG-OD2	5.67	123.40	118.30
1	B	65	ASP	CB-CG-OD2	5.67	123.40	118.30
1	B	304	ASP	CB-CG-OD2	5.56	123.30	118.30
1	A	267	ASP	CB-CG-OD2	5.43	123.19	118.30
1	A	182	ASP	CB-CG-OD2	5.43	123.19	118.30
1	A	38	ASP	CB-CG-OD2	5.22	123.00	118.30
1	B	125	ASP	CB-CG-OD2	5.14	122.93	118.30
1	A	254	ASP	CB-CG-OD2	5.09	122.88	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2593	0	2518	26	0
1	B	2617	0	2554	47	0
2	A	43	0	32	0	0
2	B	43	0	32	1	0
3	A	4	0	6	1	0
3	B	4	0	6	4	0
4	A	14	0	13	0	0
4	B	14	0	13	5	0
5	A	175	0	0	3	1
5	B	199	0	0	9	1
All	All	5706	0	5174	79	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:TYR:H	3:B:343:DMS:H22	1.07	1.12
1:B:257:VAL:CG2	5:B:369:HOH:O	1.99	1.11
4:B:344:NAG:H82	4:B:344:NAG:H3	1.39	1.02
1:B:139:ARG:HG2	1:B:139:ARG:HH11	1.24	1.01
4:B:344:NAG:C8	4:B:344:NAG:H3	1.91	0.98
1:B:20:TYR:H	3:B:343:DMS:C2	1.77	0.96
1:B:20:TYR:N	3:B:343:DMS:H22	1.88	0.88
1:B:14:ASN:HD21	1:B:17:ASP:H	1.26	0.82
1:B:71:SER:HB2	1:B:96:ILE:HD13	1.63	0.80
1:A:14:ASN:HD21	1:A:17:ASP:H	1.30	0.77
1:B:139:ARG:HH11	1:B:139:ARG:CG	1.96	0.76
4:B:344:NAG:C8	4:B:344:NAG:C3	2.65	0.75
1:B:196:ILE:HD11	1:B:204:GLN:HG2	1.74	0.69
1:A:48:SER:HB2	1:A:110:GLU:HB3	1.75	0.67
1:B:14:ASN:HD22	1:B:14:ASN:C	1.98	0.67
1:B:26:ILE:HG12	1:B:97:ILE:HG12	1.80	0.64
1:B:48:SER:HB2	1:B:110:GLU:HB3	1.80	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:ARG:HG2	1:B:139:ARG:NH1	2.02	0.63
1:B:206:LYS:HE3	1:B:274:GLY:H	1.63	0.63
1:B:257:VAL:HG23	5:B:369:HOH:O	1.76	0.62
1:A:251:ARG:HB2	1:A:254:ASP:OD1	2.01	0.61
1:A:14:ASN:ND2	1:A:17:ASP:H	1.99	0.60
1:A:198:THR:HG22	1:A:332:ASN:CG	2.23	0.59
1:B:221:CYS:HB2	5:B:402:HOH:O	2.02	0.59
1:A:14:ASN:HD21	1:A:17:ASP:N	1.99	0.57
1:B:196:ILE:CD1	1:B:204:GLN:HG2	2.33	0.57
1:B:14:ASN:ND2	1:B:17:ASP:H	2.00	0.57
1:B:218:GLU:O	1:B:219:ASP:HB2	2.04	0.56
1:B:14:ASN:HD21	1:B:17:ASP:N	2.00	0.56
1:B:275:LYS:HD3	1:B:277:TYR:CZ	2.42	0.55
1:A:164:ARG:HD2	5:A:506:HOH:O	2.07	0.55
1:B:257:VAL:HG21	5:B:369:HOH:O	1.87	0.54
1:B:96:ILE:HD12	1:B:96:ILE:N	2.23	0.54
1:B:71:SER:CB	1:B:96:ILE:HD13	2.35	0.54
1:B:341:HIS:CE1	5:B:491:HOH:O	2.60	0.53
1:B:79:LEU:HD21	1:B:92:LEU:HG	1.91	0.53
1:B:209:SER:HB2	1:B:269:SER:HB2	1.93	0.50
1:B:71:SER:HB2	1:B:96:ILE:CD1	2.38	0.50
1:A:79:LEU:HD11	1:A:92:LEU:HD23	1.94	0.50
1:B:14:ASN:C	1:B:14:ASN:ND2	2.65	0.50
1:A:206:LYS:NZ	1:A:273:GLY:O	2.45	0.49
1:A:285:VAL:HG22	1:A:298:LEU:HD22	1.95	0.48
1:B:286:PHE:HD2	1:B:297:THR:HG23	1.77	0.48
1:B:139:ARG:HG3	5:B:385:HOH:O	2.13	0.48
1:A:74[A]:HIS:CD2	5:A:472:HOH:O	2.67	0.47
1:A:16:MET:O	1:A:17:ASP:HB2	2.14	0.47
1:B:185:HIS:HD2	5:B:468:HOH:O	1.98	0.46
4:B:344:NAG:H83	4:B:344:NAG:C3	2.43	0.46
1:A:160:PHE:CZ	1:A:176:ILE:HD12	2.51	0.46
1:B:139:ARG:CG	1:B:139:ARG:NH1	2.66	0.46
1:A:259:CYS:HB2	1:A:291:SER:O	2.16	0.46
1:A:18:THR:O	3:A:343:DMS:H11	2.16	0.46
1:B:163:ASN:HD22	1:B:164:ARG:H	1.63	0.46
1:B:36:VAL:HG22	3:B:343:DMS:H21	1.98	0.45
1:B:161:TYR:HB2	1:B:324:TYR:CE2	2.51	0.45
1:B:208:VAL:HB	1:B:216:LEU:HB2	1.98	0.45
1:A:14:ASN:HD22	1:A:14:ASN:C	2.20	0.45
1:B:187:GLU:HB3	1:B:340:ARG:HD3	1.98	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:LEU:HD11	1:A:268:ILE:HD11	2.00	0.44
1:B:8:SER:HB2	5:B:511:HOH:O	2.17	0.44
1:A:80:THR:HG23	1:A:87:THR:HG23	1.99	0.44
1:A:232:ILE:HG13	1:A:318:THR:HB	1.99	0.43
1:B:12:LEU:HD21	1:B:176[A]:ILE:HD12	2.00	0.43
1:A:80:THR:HG23	1:A:87:THR:CG2	2.49	0.43
1:A:4:GLY:HA2	5:A:385:HOH:O	2.18	0.43
1:A:8:SER:OG	1:A:100:GLY:O	2.28	0.42
1:A:128:VAL:HG13	1:A:143:ILE:HG21	2.01	0.42
1:B:2:THR:HG23	1:B:4:GLY:H	1.85	0.42
1:A:305:ILE:HA	1:A:306:PRO:HD3	1.79	0.42
1:B:11:ILE:HD12	1:B:172:LEU:HD11	2.02	0.42
1:B:16:MET:O	1:B:17:ASP:HB2	2.20	0.42
2:B:342:A5T:H31	2:B:342:A5T:C32	2.49	0.42
1:B:303:MET:HG3	5:B:534:HOH:O	2.19	0.41
4:B:344:NAG:H83	4:B:344:NAG:H3	1.89	0.41
1:A:196:ILE:HD11	1:A:204:GLN:N	2.36	0.41
1:B:56:THR:HA	1:B:59:VAL:HG12	2.02	0.41
1:B:117:LEU:HA	1:B:118:PRO:HA	1.86	0.41
1:B:218:GLU:O	1:B:219:ASP:CB	2.70	0.40
1:A:161:TYR:HB2	1:A:324:TYR:CE2	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:370:HOH:O	5:B:371:HOH:O[4_456]	2.02	0.18

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/341 (97%)	326 (98%)	6 (2%)	0	100	100
1	B	335/341 (98%)	330 (98%)	5 (2%)	0	100	100
All	All	667/682 (98%)	656 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/291 (98%)	276 (96%)	10 (4%)	36	35
1	B	289/291 (99%)	271 (94%)	18 (6%)	18	13
All	All	575/582 (99%)	547 (95%)	28 (5%)	25	21

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ASN
1	A	81	LEU
1	A	135	GLN
1	A	139	ARG
1	A	215	LEU
1	A	219	ASP
1	A	251	ARG
1	A	293	LYS
1	A	309	THR
1	A	333	ARG
1	B	2	THR
1	B	6	THR
1	B	14	ASN
1	B	62	LYS
1	B	80	THR
1	B	81	LEU
1	B	87	THR
1	B	88	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	104	VAL
1	B	139	ARG
1	B	140	VAL
1	B	145	ASP
1	B	152	VAL
1	B	163	ASN
1	B	166	SER
1	B	206	LYS
1	B	288	GLU
1	B	330	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	ASN
1	A	94	GLN
1	A	150	GLN
1	A	194	ASN
1	A	341	HIS
1	B	14	ASN
1	B	74	HIS
1	B	135	GLN
1	B	163	ASN
1	B	185	HIS
1	B	194	ASN
1	B	202	GLN
1	B	287	GLN
1	B	331	ASN
1	B	341	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	DMS	A	343	-	3,3,3	2.53	1 (33%)	3,3,3	0.62	0
2	A5T	B	342	-	45,48,48	1.30	6 (13%)	59,69,69	1.64	6 (10%)
4	NAG	A	344	1	14,14,15	0.50	0	17,19,21	1.18	2 (11%)
4	NAG	B	344	1	14,14,15	1.02	1 (7%)	17,19,21	2.15	5 (29%)
2	A5T	A	342	-	45,48,48	1.36	7 (15%)	59,69,69	1.73	11 (18%)
3	DMS	B	343	-	3,3,3	2.83	1 (33%)	3,3,3	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	344	1	-	4/6/23/26	0/1/1/1
4	NAG	B	344	1	-	5/6/23/26	0/1/1/1
2	A5T	A	342	-	-	5/27/51/51	0/6/6/6
2	A5T	B	342	-	-	1/27/51/51	0/6/6/6

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	343	DMS	O-S	4.84	1.82	1.50
3	A	343	DMS	O-S	4.28	1.79	1.50
2	B	342	A5T	C10-C12	-3.74	1.34	1.52

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	342	A5T	C2-CL11	3.73	1.80	1.72
2	A	342	A5T	C10-C12	-3.65	1.34	1.52
2	A	342	A5T	C2-CL11	3.45	1.80	1.72
2	B	342	A5T	C33-CLR3	3.24	1.79	1.72
2	A	342	A5T	C24-N27	-3.14	1.45	1.47
2	A	342	A5T	C33-CLR3	2.96	1.79	1.72
2	A	342	A5T	C21-C19	2.95	1.54	1.50
4	B	344	NAG	O5-C1	-2.68	1.39	1.43
2	B	342	A5T	C24-N27	-2.60	1.45	1.47
2	B	342	A5T	C36-CLA	2.58	1.79	1.73
2	A	342	A5T	C36-CLA	2.50	1.79	1.73
2	A	342	A5T	C31-C28	2.20	1.53	1.48
2	B	342	A5T	C30-C28	2.08	1.53	1.48

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	342	A5T	C29-N26-C28	-6.39	111.65	118.33
2	B	342	A5T	C20-C22-N26	5.36	125.23	119.66
2	B	342	A5T	C29-N26-C28	-5.36	112.73	118.33
4	B	344	NAG	C2-N2-C7	4.81	129.76	122.90
4	B	344	NAG	C1-O5-C5	3.75	117.27	112.19
4	B	344	NAG	O5-C5-C6	3.67	112.96	107.20
2	A	342	A5T	C3-C2-C4	3.58	120.97	117.93
2	A	342	A5T	C32-C29-N26	3.48	118.32	113.34
2	A	342	A5T	C34-C32-C33	3.44	119.87	117.43
2	B	342	A5T	C33-C36-CLA	-3.44	117.17	120.52
2	A	342	A5T	C20-C22-N26	3.43	123.22	119.66
2	B	342	A5T	C32-C29-N26	3.30	118.06	113.34
4	A	344	NAG	O5-C5-C6	3.23	112.27	107.20
2	B	342	A5T	C29-C32-C33	-2.73	117.19	121.73
2	A	342	A5T	C29-C32-C33	-2.70	117.25	121.73
2	A	342	A5T	C12-C10-C8	2.62	122.16	113.34
4	B	344	NAG	O3-C3-C2	-2.60	104.08	109.47
4	A	344	NAG	O5-C1-C2	-2.55	107.26	111.29
2	A	342	A5T	F1-C6-C3	2.48	120.97	117.58
2	A	342	A5T	C24-C21-C19	2.35	114.20	110.37
2	A	342	A5T	O5-C3-C2	-2.34	117.55	120.66
2	B	342	A5T	C24-C21-C19	2.23	114.00	110.37
4	B	344	NAG	C6-C5-C4	-2.21	107.83	113.00
2	A	342	A5T	C20-C23-N27	2.16	111.60	108.91

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	344	NAG	C8-C7-N2-C2
4	A	344	NAG	O7-C7-N2-C2
4	B	344	NAG	C3-C2-N2-C7
2	A	342	A5T	C2-C3-O5-C8
4	A	344	NAG	O5-C5-C6-O6
4	B	344	NAG	C4-C5-C6-O6
4	A	344	NAG	C4-C5-C6-O6
4	B	344	NAG	C8-C7-N2-C2
4	B	344	NAG	O7-C7-N2-C2
4	B	344	NAG	O5-C5-C6-O6
2	A	342	A5T	C12-C10-C8-O5
2	A	342	A5T	C8-C10-C12-C13
2	A	342	A5T	C10-C8-O5-C3
2	A	342	A5T	C31-C28-N26-C22
2	B	342	A5T	C12-C10-C8-O5


There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	343	DMS	1	0
2	B	342	A5T	1	0
4	B	344	NAG	5	0
3	B	343	DMS	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	334/341 (97%)	0.05	10 (2%) 50 49	19, 33, 56, 70	0
1	B	337/341 (98%)	0.15	10 (2%) 50 49	19, 35, 53, 72	0
All	All	671/682 (98%)	0.10	20 (2%) 50 49	19, 34, 55, 72	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	SER	5.0
1	A	213	SER	4.6
1	A	252	LEU	3.1
1	A	214	THR	2.7
1	B	69	SER	2.6
1	B	219	ASP	2.5
1	B	290	TYR	2.5
1	A	219	ASP	2.4
1	B	259	CYS	2.4
1	B	292	SER	2.4
1	A	264	THR	2.3
1	A	293	LYS	2.3
1	B	253	PHE	2.3
1	A	239	ILE	2.3
1	A	294	LYS	2.2
1	B	293	LYS	2.2
1	A	216	LEU	2.2
1	B	53	ARG	2.2
1	B	260	ASN	2.2
1	B	294	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	B	344	14/15	0.69	0.30	64,69,72,72	0
4	NAG	A	344	14/15	0.76	0.25	51,56,59,59	0
2	A5T	B	342	43/43	0.91	0.15	25,33,42,44	0
2	A5T	A	342	43/43	0.93	0.12	25,32,40,43	0
3	DMS	A	343	4/4	0.95	0.24	40,42,44,46	0
3	DMS	B	343	4/4	0.98	0.15	34,37,37,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around A5T B 342:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.