



Full wwPDB X-ray Structure Validation Report i

Nov 5, 2023 – 10:32 pm GMT

PDB ID : 6HQF
Title : Structure of Phenylalanine ammonia-lyase from Petroselinum crispum in complex with (R)-APEP
Authors : Bata, Z.; Molnar, B.; Leveles, I.; Poppe, L.; Vertessy, G.B.
Deposited on : 2018-09-24
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 10955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phenylalanine ammonia-lyase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	673	5142	3246	884	989	23	0	4	0
1	B	673	5135	3243	880	988	24	0	2	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	MDO	ALA	chromophore	UNP P24481
A	?	MDO	SER	chromophore	UNP P24481
A	203	MDO	GLY	chromophore	UNP P24481
A	704	SER	CYS	engineered mutation	UNP P24481
A	716	SER	CYS	engineered mutation	UNP P24481
B	?	MDO	ALA	chromophore	UNP P24481
B	?	MDO	SER	chromophore	UNP P24481
B	203	MDO	GLY	chromophore	UNP P24481
B	704	SER	CYS	engineered mutation	UNP P24481
B	716	SER	CYS	engineered mutation	UNP P24481

- Molecule 2 is [(1R)-1-amino-2-phenylethyl]phosphonic acid (three-letter code: PPH) (formula: C₈H₁₂NO₃P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total C N O P					0	0
			13 8 1 3 1						
2	B	1	Total C N O P					0	0
			13 8 1 3 1						


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	327	Total O 334 334		0	7
3	B	310	Total O 318 318		0	8


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phenylalanine ammonia-lyase 1



- Molecule 1: Phenylalanine ammonia-lyase 1



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	118.88Å 160.95Å 141.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.70 – 1.76 46.22 – 1.72	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.70-1.76) 98.6 (46.22-1.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	0.95 (at 1.72Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R , R_{free}	0.168 , 0.205 0.168 , 0.204	Depositor DCC
R_{free} test set	2098 reflections (1.49%)	wwPDB-VP
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.078	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.4	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10955	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Continued from previous page...

Mol	Chain	Res	Type
1	A	260	ASN
1	A	484	GLU
1	A	560	ARG
1	A	562	CYS
1	A	575	ILE
1	A	612	SER
1	A	665	ARG
1	B	25	GLU
1	B	67	SER
1	B	139	ASN
1	B	260	ASN
1	B	382	ASN
1	B	484	GLU
1	B	547	VAL
1	B	549	THR
1	B	562	CYS
1	B	575	ILE
1	B	674	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	260	ASN
1	B	260	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

5.8 Polymer linkage issues

There are no chain breaks in this entry.

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	612	SER	2.8
1	B	643	SER	2.8
1	B	475	PRO	2.8
1	A	476	VAL	2.8
1	A	407	VAL	2.7
1	B	560	ARG	2.7
1	B	469	LEU	2.7
1	B	543	VAL	2.7
1	A	612	SER	2.7
1	B	78	SER	2.7
1	A	108	ASP	2.7
1	B	424	PHE	2.7
1	B	422	LEU	2.6
1	B	602	LEU	2.6
1	B	112	VAL	2.6
1	B	366	VAL	2.6
1	B	466	CYS	2.6
1	A	425	ALA	2.6
1	B	121	HIS	2.6
1	A	493	LEU	2.6
1	B	472	LEU	2.6
1	B	351	TYR	2.6
1	A	347	LYS	2.6
1	B	418	ALA	2.6
1	B	629	LEU	2.5
1	B	110	TYR	2.5
1	B	477	THR	2.5
1	A	632	LYS	2.5
1	B	76	ASP	2.5
1	A	25	GLU	2.5
1	B	603	LYS	2.5
1	A	32	ILE	2.5
1	B	547	VAL	2.5
1	A	123	ARG	2.4
1	B	415	ALA	2.4
1	A	463	ALA	2.4
1	B	478	ASN	2.4
1	B	200	ILE	2.4
1	A	121	HIS	2.4
1	A	244	GLU	2.4
1	A	355	THR	2.4
1	B	80	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	508	LEU	2.3
1	B	123	ARG	2.3
1	B	479	HIS	2.3
1	A	501	THR	2.3
1	B	614	SER	2.3
1	A	120	SER	2.3
1	A	231	ILE	2.3
1	A	479	HIS	2.3
1	B	419	ILE	2.3
1	B	604	ASN	2.3
1	A	421	LYS	2.3
1	B	111	GLY	2.3
1	B	124	THR	2.3
1	B	605	GLY	2.2
1	B	407	VAL	2.2
1	A	111	GLY	2.2
1	B	406	GLY	2.2
1	B	617	GLN	2.2
1	A	77	GLY	2.2
1	B	403	THR	2.2
1	A	496	ILE	2.2
1	A	359	TRP	2.2
1	A	497	SER	2.1
1	A	510	LEU	2.1
1	A	641	LEU	2.1
1	A	351	TYR	2.1
1	A	545	LYS	2.1
1	A	366	VAL	2.1
1	A	460	ILE	2.1
1	B	606	ASP	2.1
1	B	483	ALA	2.1
1	B	474	ASN	2.1
1	A	559	SER	2.1
1	B	228	THR	2.0
1	A	480	VAL	2.0
1	B	548	LEU	2.0
1	B	632	LYS	2.0
1	B	551	GLY	2.0
1	A	461	ALA	2.0
1	B	461	ALA	2.0
1	A	465	TYR	2.0
1	B	101	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MDO	B	203	13/14	0.92	0.12	20,24,31,34	0
1	MDO	A	203	13/14	0.93	0.12	22,24,32,34	0

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PPH	A	801	13/13	0.96	0.08	23,28,33,35	0
2	PPH	B	801	13/13	0.97	0.09	25,30,35,36	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.