wwPDB X-ray Structure Validation Summary Report (i)

Aug 27, 2023 – 02:41 PM EDT

PDB ID	:	3HZG
Title	:	Crystal structure of mycobacterium tuberculosis thymidylate synthase X bound with FAD
Authors	:	Staker, B.L.; Rathod, P.; Hunter, J.; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on	:	2009-06-23
Resolution	:	2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	6% 79% 15% 5%
1	B	258	9% 76% 18% 6%
1	C	258	2% 78% 17% 5%
1	D	258	5% 75% 19% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	259	-	-	X	-
4	GOL	C	259	-	-	-	X

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8129 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thymidylate synthase thyX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	1911	1203	349	354	5	0	0	0
1	B	243	1889	1188	345	351	5	0	1	0
1	C	245	1904	1199	348	352	5	0	0	0
1	D	245	1893	1191	343	354	5	0	0	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	251	LEU	-	expression tag	UNP P66930
A	252	GLU	-	expression tag	UNP P66930
A	253	HIS	-	expression tag	UNP P66930
A	254	HIS	-	expression tag	UNP P66930
A	255	HIS	-	expression tag	UNP P66930
A	256	HIS	-	expression tag	UNP P66930
A	257	HIS	-	expression tag	UNP P66930
A	258	HIS	-	expression tag	UNP P66930
B	251	LEU	-	expression tag	UNP P66930
B	252	GLU	-	expression tag	UNP P66930
B	253	HIS	-	expression tag	UNP P66930
B	254	HIS	-	expression tag	UNP P66930
B	255	HIS	-	expression tag	UNP P66930
B	256	HIS	-	expression tag	UNP P66930
B	257	HIS	-	expression tag	UNP P66930
B	258	HIS	-	expression tag	UNP P66930
C	251	LEU	-	expression tag	UNP P66930
C	252	GLU	-	expression tag	UNP P66930
C	253	HIS	-	expression tag	UNP P66930
C	254	HIS	-	expression tag	UNP P66930
C	255	HIS	-	expression tag	UNP P66930

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	256	HIS	-	expression tag	UNP P66930
C	257	HIS	-	expression tag	UNP P66930
C	258	HIS	-	expression tag	UNP P66930
D	251	LEU	-	expression tag	UNP P66930
D	252	GLU	-	expression tag	UNP P66930
D	253	HIS	-	expression tag	UNP P66930
D	254	HIS	-	expression tag	UNP P66930
D	255	HIS	-	expression tag	UNP P66930
D	256	HIS	-	expression tag	UNP P66930
D	257	HIS	-	expression tag	UNP P66930
D	258	HIS	-	expression tag	UNP P66930

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	O	P			
2	A	1	5	4	1		0	0
2	A	1	5	4	1		0	0
2	B	1	5	4	1		0	0
2	B	1	5	4	1		0	0
2	B	1	5	4	1		0	0
2	C	1	5	4	1		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	C	1	5	4	1	0	0
			Total	O	P		
2	D	1	5	4	1	0	0
			Total	O	P		
2	D	1	5	4	1	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	53	27	9	15	2	0	0
			Total	C	N	O	P		
3	B	1	53	27	9	15	2	0	0
			Total	C	N	O	P		
3	C	1	53	27	9	15	2	0	0
			Total	C	N	O	P		
3	D	1	53	27	9	15	2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0
			Total	C	O		
4	C	1	6	3	3	0	0
			Total	C	O		
4	C	1	6	3	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	64	64	64	0	0
			Total	O		
5	B	58	58	58	0	0
			Total	O		
5	C	76	76	76	0	0
			Total	O		
5	D	59	59	59	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thymidylate synthase thyX

Chain A:



• Molecule 1: Thymidylate synthase thyX

Chain B:



• Molecule 1: Thymidylate synthase thyX

Chain C:



• Molecule 1: Thymidylate synthase thyX

Chain D:


MET
A2
L7
A13
D16
V36
Y44
P50
N51
P52
I65
V72
S79
I82
T83
R87
T90
I94
R95
Q103
L104
S105
Y108
V109
P110
E111
R115
V116
M122
T142
E145
K149
L150
E151
A152
K153
F154
A155
D156
Q157
P158
N159
A160
I161
L162
R163
R168
Q169
R172
A173
V174
L175
E180
T181
R182
I183
H194
R199
A200
S201
L211
A212
I213
E231
A242
P245
L246
ALA
THR
GLU
ALA
LEU
GLU
HIS
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	78.55Å 86.59Å 162.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.30 – 2.45 43.30 – 2.45	Depositor EDS
% Data completeness (in resolution range)	(Not available) (43.30-2.45) 99.6 (43.30-2.45)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.63 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.198 , 0.273 0.196 , 0.269	Depositor DCC
R_{free} test set	2081 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å^2)	39.1	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 45.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	8129	wwPDB-VP
Average B, all atoms (Å^2)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, PO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.72	0/1955	0.76	0/2664
1	B	0.73	0/1937	0.76	1/2640 (0.0%)
1	C	0.77	0/1948	0.74	0/2656
1	D	0.74	0/1937	0.75	1/2645 (0.0%)
All	All	0.74	0/7777	0.76	2/10605 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	199	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	B	18	LEU	CA-CB-CG	5.22	127.31	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1911	0	1891	23	0
1	B	1889	0	1856	28	0
1	C	1904	0	1878	24	0
1	D	1893	0	1847	45	0
2	A	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	15	0	0	0	0
2	C	10	0	0	1	0
2	D	10	0	0	0	0
3	A	53	0	31	2	0
3	B	53	0	31	0	0
3	C	53	0	31	0	0
3	D	53	0	31	4	0
4	A	6	0	8	4	0
4	C	12	0	16	0	0
5	A	64	0	0	0	0
5	B	58	0	0	0	0
5	C	76	0	0	1	0
5	D	59	0	0	2	0
All	All	8129	0	7620	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:36:VAL:HG13	1:C:72:VAL:HG12	1.53	0.89
1:C:13:ALA:HB3	1:C:79:SER:HB2	1.60	0.82
1:B:51:ASN:O	1:B:53:LYS:N	2.23	0.70
1:C:36:VAL:HG13	1:C:72:VAL:CG1	2.22	0.69
1:D:13:ALA:HB3	1:D:79:SER:HB2	1.72	0.69

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/258 (94%)	228 (94%)	15 (6%)	0	100	100
1	B	242/258 (94%)	232 (96%)	7 (3%)	3 (1%)	13	12
1	C	243/258 (94%)	236 (97%)	6 (2%)	1 (0%)	34	41
1	D	243/258 (94%)	234 (96%)	9 (4%)	0	100	100
All	All	971/1032 (94%)	930 (96%)	37 (4%)	4 (0%)	34	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	52	PRO
1	C	236	ALA
1	B	49	LYS
1	B	24	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/209 (95%)	186 (94%)	12 (6%)	18	24
1	B	194/209 (93%)	190 (98%)	4 (2%)	53	66
1	C	196/209 (94%)	185 (94%)	11 (6%)	21	27
1	D	194/209 (93%)	184 (95%)	10 (5%)	23	30
All	All	782/836 (94%)	745 (95%)	37 (5%)	26	34

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	72	VAL
1	D	201	SER
1	D	116	VAL
1	D	161	ILE
1	B	162	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such

sidechains are listed below:

Mol	Chain	Res	Type
1	D	157	GLN
1	D	159	ASN
1	D	169	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	PO4	B	1007	-	4,4,4	0.67	0	6,6,6	0.76	0
2	PO4	C	1004	-	4,4,4	0.76	0	6,6,6	0.89	0
2	PO4	C	1006	-	4,4,4	0.82	0	6,6,6	0.94	0
3	FAD	C	501	-	53,58,58	1.54	6 (11%)	68,89,89	1.98	18 (26%)
4	GOL	C	259	-	5,5,5	0.34	0	5,5,5	0.29	0
3	FAD	D	501	-	53,58,58	1.43	8 (15%)	68,89,89	1.97	19 (27%)
2	PO4	A	1002	-	4,4,4	0.91	0	6,6,6	0.70	0
2	PO4	D	1003	-	4,4,4	0.87	0	6,6,6	0.95	0
3	FAD	A	501	-	53,58,58	1.32	6 (11%)	68,89,89	1.55	14 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	A	259	-	5,5,5	0.47	0	5,5,5	0.72	0
2	PO4	B	1009	-	4,4,4	1.05	0	6,6,6	1.38	1 (16%)
2	PO4	A	1001	-	4,4,4	1.33	0	6,6,6	0.83	0
3	FAD	B	501	-	53,58,58	1.17	3 (5%)	68,89,89	1.60	16 (23%)
2	PO4	D	1008	-	4,4,4	0.81	0	6,6,6	0.44	0
2	PO4	B	1005	-	4,4,4	1.40	0	6,6,6	0.76	0
4	GOL	C	260	-	5,5,5	0.43	0	5,5,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	C	501	-	-	9/30/50/50	0/6/6/6
3	FAD	D	501	-	-	10/30/50/50	0/6/6/6
3	FAD	A	501	-	-	11/30/50/50	0/6/6/6
4	GOL	A	259	-	-	0/4/4/4	-
4	GOL	C	259	-	-	2/4/4/4	-
3	FAD	B	501	-	-	10/30/50/50	0/6/6/6
4	GOL	C	260	-	-	2/4/4/4	-

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	501	FAD	C1'-C2'	5.87	1.60	1.52
3	B	501	FAD	C4X-N5	4.67	1.39	1.30
3	D	501	FAD	C1'-C2'	4.41	1.58	1.52
3	D	501	FAD	C4X-N5	3.73	1.38	1.30
3	C	501	FAD	C4X-N5	3.65	1.37	1.30

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	501	FAD	C1'-C2'-C3'	5.59	125.42	109.79
3	B	501	FAD	N3A-C2A-N1A	-5.49	120.10	128.68
3	C	501	FAD	O3'-C3'-C2'	5.46	122.01	108.81
3	D	501	FAD	N3A-C2A-N1A	-5.33	120.34	128.68
3	D	501	FAD	O3'-C3'-C2'	5.32	121.66	108.81

There are no chirality outliers.

5 of 44 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	FAD	C5B-O5B-PA-O1A
3	A	501	FAD	C1'-C2'-C3'-O3'
3	A	501	FAD	C1'-C2'-C3'-C4'
3	A	501	FAD	O2'-C2'-C3'-O3'
3	A	501	FAD	O2'-C2'-C3'-C4'

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1006	PO4	1	0
3	D	501	FAD	4	0
3	A	501	FAD	2	0
4	A	259	GOL	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FAD C 501



Bond lengths



Bond angles



Torsions



Rings

Ligand FAD D 501



Bond lengths



Bond angles



Torsions



Rings

Ligand FAD A 501



Bond lengths



Bond angles



Torsions



Rings

Ligand FAD B 501



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	245/258 (94%)	0.28	16 (6%)	18	15	21, 36, 59, 66	0
1	B	243/258 (94%)	0.35	22 (9%)	9	6	18, 35, 56, 74	1 (0%)
1	C	245/258 (94%)	-0.07	5 (2%)	65	62	15, 29, 52, 64	0
1	D	245/258 (94%)	0.17	14 (5%)	23	20	18, 33, 64, 71	0
All	All	978/1032 (94%)	0.18	57 (5%)	23	20	15, 33, 59, 74	1 (0%)

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	236	ALA	4.6
1	A	246	LEU	4.5
1	D	161	ILE	4.3
1	C	237	ASP	4.1
1	B	50	PRO	3.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	C	259	6/6	0.70	0.54	66,67,68,69	0
4	GOL	A	259	6/6	0.78	0.38	46,50,51,52	0
2	PO4	B	1007	5/5	0.91	0.21	60,60,61,62	0
2	PO4	A	1002	5/5	0.92	0.15	76,77,78,78	0
4	GOL	C	260	6/6	0.93	0.30	55,57,58,58	0
2	PO4	B	1009	5/5	0.94	0.22	60,60,62,63	0
3	FAD	A	501	53/53	0.94	0.18	26,37,49,50	0
3	FAD	B	501	53/53	0.94	0.18	26,39,62,63	0
2	PO4	D	1008	5/5	0.95	0.10	50,52,55,55	0
3	FAD	D	501	53/53	0.96	0.16	18,25,32,36	0
3	FAD	C	501	53/53	0.97	0.19	16,25,28,33	0
2	PO4	C	1004	5/5	0.98	0.20	37,37,38,39	0
2	PO4	D	1003	5/5	0.98	0.16	35,36,37,40	0
2	PO4	C	1006	5/5	0.99	0.12	28,28,30,30	0
2	PO4	B	1005	5/5	0.99	0.17	26,27,28,30	0
2	PO4	A	1001	5/5	1.00	0.15	24,24,26,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 501:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around FAD B 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FAD D 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FAD C 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.