



Full wwPDB NMR Structure Validation Report ⓘ

May 29, 2020 – 07:13 am BST

PDB ID : 5IX9
Title : Cell surface anchoring domain
Authors : Guo, S.; Langelaan, D.
Deposited on : 2016-03-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 92%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	195	

2 Ensemble composition and analysis

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:41, A:45-A:71 (67)	0.15	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	3, 6, 8, 9, 10, 14, 17, 19
2	1, 2, 13, 16, 18
3	4, 11, 15, 20
Single-model clusters	5; 7; 12

3 Entry composition



There is only 1 type of molecule in this entry. The entry contains 1084 atoms, of which 535 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Antifreeze protein.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	76	1084	343	535	87	118	1	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	initiating methionine	UNP A1YIY2
A	-9	SER	-	expression tag	UNP A1YIY2
A	-8	ASP	-	expression tag	UNP A1YIY2
A	-7	ASN	-	expression tag	UNP A1YIY2
A	-6	GLN	-	expression tag	UNP A1YIY2
A	-5	PHE	-	expression tag	UNP A1YIY2
A	-4	PRO	-	expression tag	UNP A1YIY2
A	-3	PHE	-	expression tag	UNP A1YIY2
A	-2	ALA	-	expression tag	UNP A1YIY2
A	-1	THR	-	expression tag	UNP A1YIY2
A	0	LEU	-	expression tag	UNP A1YIY2
A	20	ASN	ASP	engineered mutation	UNP A1YIY2
A	177	LEU	-	expression tag	UNP A1YIY2
A	178	GLU	-	expression tag	UNP A1YIY2
A	179	HIS	-	expression tag	UNP A1YIY2
A	180	HIS	-	expression tag	UNP A1YIY2
A	181	HIS	-	expression tag	UNP A1YIY2
A	182	HIS	-	expression tag	UNP A1YIY2
A	183	HIS	-	expression tag	UNP A1YIY2
A	184	HIS	-	expression tag	UNP A1YIY2



4.2.9 Score per residue for model 9

- Molecule 1: Antifreeze protein



4.2.10 Score per residue for model 10

- Molecule 1: Antifreeze protein



4.2.11 Score per residue for model 11

- Molecule 1: Antifreeze protein



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *na*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	refinement	
CNS	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	889
Number of shifts mapped to atoms	889
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	92%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	496	495	495	11±2
All	All	9920	9900	9900	218

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:4:ILE:HB	1:A:71:TYR:CE2	0.73	2.19	13	20
1:A:4:ILE:HB	1:A:71:TYR:CD2	0.64	2.28	19	20
1:A:52:ASP:HB2	1:A:71:TYR:OH	0.63	1.93	7	4
1:A:45:SER:HA	1:A:58:ILE:O	0.59	1.98	7	8
1:A:16:VAL:O	1:A:23:GLU:HA	0.59	1.98	11	15
1:A:16:VAL:HG12	1:A:24:ARG:HG2	0.55	1.79	3	13
1:A:38:VAL:O	1:A:63:ILE:HA	0.55	2.00	10	9
1:A:66:MET:SD	1:A:70:ILE:HG21	0.53	2.43	12	4
1:A:9:LYS:HE2	1:A:47:THR:OG1	0.53	2.03	9	2
1:A:50:PHE:CE1	1:A:56:VAL:HB	0.52	2.40	19	7
1:A:46:VAL:O	1:A:57:VAL:HA	0.51	2.04	9	14
1:A:27:LYS:O	1:A:30:ASP:HB3	0.50	2.06	2	2
1:A:26:LEU:HD22	1:A:30:ASP:OD1	0.50	2.07	13	9
1:A:7:ILE:HG12	1:A:30:ASP:O	0.49	2.08	9	10
1:A:47:THR:HA	1:A:56:VAL:O	0.49	2.08	1	5
1:A:50:PHE:HB3	1:A:71:TYR:CE1	0.47	2.45	4	5
1:A:66:MET:HE3	1:A:70:ILE:HG21	0.47	1.86	6	2
1:A:10:LEU:HD11	1:A:14:VAL:HG21	0.46	1.87	2	8
1:A:9:LYS:HB3	1:A:47:THR:HB	0.46	1.87	16	10
1:A:66:MET:CE	1:A:70:ILE:HG21	0.46	2.41	5	3
1:A:48:ILE:HB	1:A:56:VAL:CG2	0.44	2.42	10	11
1:A:36:GLU:O	1:A:66:MET:HG2	0.44	2.12	11	5
1:A:26:LEU:HD22	1:A:30:ASP:CG	0.43	2.33	2	7
1:A:9:LYS:HB2	1:A:47:THR:HB	0.43	1.90	4	1
1:A:14:VAL:HG23	1:A:26:LEU:HB2	0.43	1.89	15	6
1:A:48:ILE:HD13	1:A:66:MET:HE2	0.43	1.91	8	2
1:A:50:PHE:CE1	1:A:56:VAL:CG2	0.43	3.02	8	6
1:A:7:ILE:HG13	1:A:30:ASP:OD1	0.43	2.14	1	1
1:A:16:VAL:HG22	1:A:36:GLU:HB3	0.42	1.91	14	3
1:A:32:ILE:CG2	1:A:66:MET:HG3	0.42	2.44	14	1
1:A:9:LYS:HB3	1:A:47:THR:CB	0.42	2.44	16	1
1:A:22:GLN:HA	1:A:22:GLN:HE21	0.41	1.75	10	1
1:A:4:ILE:HG21	1:A:34:PHE:HD2	0.41	1.75	10	1
1:A:17:GLN:HA	1:A:22:GLN:O	0.40	2.15	5	1
1:A:24:ARG:NH2	1:A:36:GLU:OE2	0.40	2.53	12	1

5.2 Torsion angles

5.2.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	67/195 (34%)	63±1 (93±1%)	4±1 (7±1%)	0±0 (0±0%)	54	85
All	All	1340/3900 (34%)	1250 (93%)	89 (7%)	1 (0%)	54	85

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	66	MET	1

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	56/159 (35%)	51±2 (91±3%)	5±2 (9±3%)	13	60
All	All	1120/3180 (35%)	1020 (91%)	100 (9%)	13	60

All 16 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	15	THR	20
1	A	16	VAL	16
1	A	28	LEU	16
1	A	30	ASP	15
1	A	56	VAL	8
1	A	24	ARG	7
1	A	11	ASP	4
1	A	70	ILE	3
1	A	39	LEU	3
1	A	64	VAL	2
1	A	27	LYS	1
1	A	54	THR	1
1	A	65	GLU	1
1	A	45	SER	1
1	A	22	GLN	1
1	A	17	GLN	1

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

There are no ligands in this entry.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 92% for the well-defined parts and 92% for the entire structure.

6.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list*

6.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	889
Number of shifts mapped to atoms	889
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

6.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	76	-0.21 ± 0.10	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	63	-0.13 ± 0.16	None needed (< 0.5 ppm)
$^{13}\text{C}'$	75	0.10 ± 0.11	None needed (< 0.5 ppm)
^{15}N	75	0.28 ± 0.23	None needed (< 0.5 ppm)

6.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 92%, i.e. 689 atoms were assigned a chemical shift out of a possible 745. 0 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	332/333 (100%)	133/133 (100%)	133/134 (99%)	66/66 (100%)
Sidechain	313/368 (85%)	179/207 (86%)	129/152 (85%)	5/9 (56%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	44/44 (100%)	24/24 (100%)	20/20 (100%)	0/0 (—%)
Overall	689/745 (92%)	336/364 (92%)	282/306 (92%)	71/75 (95%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 92%, i.e. 750 atoms were assigned a chemical shift out of a possible 815. 0 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	373/378 (99%)	147/151 (97%)	151/152 (99%)	75/75 (100%)
Sidechain	333/393 (85%)	192/221 (87%)	134/161 (83%)	7/11 (64%)
Aromatic	44/44 (100%)	24/24 (100%)	20/20 (100%)	0/0 (—%)
Overall	750/815 (92%)	363/396 (92%)	305/333 (92%)	82/86 (95%)

6.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

6.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

