



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2023 – 03:43 PM EDT

PDB ID : 1JQB  
Title : Alcohol Dehydrogenase from Clostridium Beijerinckii: Crystal Structure of Mutant with Enhanced Thermal Stability  
Authors : Levin, I.; Frolow, F.; Bogin, O.; Peretz, M.; Hacham, Y.; Burstein, Y.  
Deposited on : 2001-08-05  
Resolution : 1.97 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	
1	B	351	
1	C	351	
1	D	351	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADP-dependent Alcohol Dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	351	2647	1679	463	483	22	0	0	0
1	B	351	2647	1679	463	483	22	0	0	0
1	C	351	2647	1679	463	483	22	0	0	0
1	D	351	2648	1679	463	484	22	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1154	THR	SER	conflict	UNP P25984
A	1165	GLU	GLN	engineered mutation	UNP P25984
A	1224	GLU	VAL	engineered mutation	UNP P25984
A	1234	LYS	GLU	conflict	UNP P25984
A	1254	LYS	SER	engineered mutation	UNP P25984
A	1304	ARG	MET	engineered mutation	UNP P25984
B	2154	THR	SER	conflict	UNP P25984
B	2165	GLU	GLN	engineered mutation	UNP P25984
B	2224	GLU	VAL	engineered mutation	UNP P25984
B	2234	LYS	GLU	conflict	UNP P25984
B	2254	LYS	SER	engineered mutation	UNP P25984
B	2304	ARG	MET	engineered mutation	UNP P25984
C	3154	THR	SER	conflict	UNP P25984
C	3165	GLU	GLN	engineered mutation	UNP P25984
C	3224	GLU	VAL	engineered mutation	UNP P25984
C	3234	LYS	GLU	conflict	UNP P25984
C	3254	LYS	SER	engineered mutation	UNP P25984
C	3304	ARG	MET	engineered mutation	UNP P25984
D	4154	THR	SER	conflict	UNP P25984
D	4165	GLU	GLN	engineered mutation	UNP P25984
D	4224	GLU	VAL	engineered mutation	UNP P25984

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	4234	LYS	GLU	conflict	UNP P25984
D	4254	LYS	SER	engineered mutation	UNP P25984
D	4304	ARG	MET	engineered mutation	UNP P25984

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0
2	C	1	Total 1	Zn 1	0	0
2	D	1	Total 1	Zn 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	109	Total 109	O 109	0	0
3	B	110	Total 110	O 110	0	0
3	C	86	Total 86	O 86	0	0
3	D	106	Total 106	O 106	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NADP-dependent Alcohol Dehydrogenase




- Molecule 1: NADP-dependent Alcohol Dehydrogenase





- Molecule 1: NADP-dependent Alcohol Dehydrogenase





● Molecule 1: NADP-dependent Alcohol Dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.70Å 147.80Å 125.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.01 – 1.97 20.01 – 1.97	Depositor EDS
% Data completeness (in resolution range)	97.9 (20.01-1.97) 98.1 (20.01-1.97)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 1.97Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.219 , 0.247 0.218 , 0.247	Depositor DCC
$R_{free}$ test set	11159 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtrriage
Anisotropy	0.306	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 47.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	11004	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/2696	0.62	0/3635
1	B	0.32	0/2696	0.62	0/3635
1	C	0.30	0/2696	0.60	0/3635
1	D	0.31	0/2697	0.62	0/3635
All	All	0.31	0/10785	0.62	0/14540

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2647	0	2685	65	0
1	B	2647	0	2685	68	0
1	C	2647	0	2685	72	0
1	D	2648	0	2685	56	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	109	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	110	0	0	0	0
3	C	86	0	0	5	0
3	D	106	0	0	2	0
All	All	11004	0	10740	242	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 242 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2102:HIS:HD2	1:B:2107:LEU:H	1.08	1.02
1:A:1102:HIS:HD2	1:A:1107:LEU:H	1.03	0.94
1:B:2200:ARG:HD2	1:B:2342:LYS:HD3	1.49	0.92
1:B:2336:LEU:HD13	1:B:2344:LEU:HD22	1.54	0.89
1:C:3102:HIS:HD2	1:C:3107:LEU:H	1.19	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	349/351 (99%)	330 (95%)	18 (5%)	1 (0%)	41 29
1	B	349/351 (99%)	333 (95%)	14 (4%)	2 (1%)	25 14
1	C	349/351 (99%)	323 (93%)	25 (7%)	1 (0%)	41 29
1	D	349/351 (99%)	329 (94%)	20 (6%)	0	100 100
All	All	1396/1404 (99%)	1315 (94%)	77 (6%)	4 (0%)	41 29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2115	PHE
1	A	1219	LYS
1	C	3044	VAL
1	B	2340	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/279 (100%)	276 (99%)	3 (1%)	73	70
1	B	279/279 (100%)	278 (100%)	1 (0%)	91	90
1	C	279/279 (100%)	272 (98%)	7 (2%)	47	39
1	D	279/279 (100%)	273 (98%)	6 (2%)	52	46
All	All	1116/1116 (100%)	1099 (98%)	17 (2%)	65	59

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	4267	TYR
1	D	4327	PHE
1	C	3143	GLU
1	C	3161	LEU
1	C	3328	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	2312	ASN
1	C	3010	ASN
1	D	4232	ASN
1	C	3232	ASN
1	D	4220	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	351/351 (100%)	0.28	20 (5%) 23 25	15, 25, 51, 59	0
1	B	351/351 (100%)	0.39	36 (10%) 6 7	12, 23, 58, 70	0
1	C	351/351 (100%)	0.71	57 (16%) 1 1	14, 29, 64, 70	0
1	D	351/351 (100%)	0.36	24 (6%) 17 19	16, 26, 51, 61	0
All	All	1404/1404 (100%)	0.44	137 (9%) 7 8	12, 26, 58, 70	0

The worst 5 of 137 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	3013	GLY	7.8
1	D	4008	GLY	7.0
1	B	2050	GLY	7.0
1	D	4218	TYR	6.9
1	B	2049	LEU	6.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	B	2353	1/1	0.90	0.06	41,41,41,41	0
2	ZN	A	1353	1/1	0.97	0.04	35,35,35,35	0
2	ZN	C	3353	1/1	0.97	0.05	41,41,41,41	0
2	ZN	D	4353	1/1	0.98	0.03	36,36,36,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.