



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 10:38 PM EDT

PDB ID : 5LWQ
Title : CeuE (H227L variant) a periplasmic protein from *Campylobacter jejuni*
Authors : Wilde, E.J.; Blagova, E.; Hughes, A.; Raines, D.J.; Moroz, O.V.; Turkenburg, J.P.; Duhme-Klair, A.-K.; Wilson, K.S.
Deposited on : 2016-09-19
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	291	<div> <div></div> <div>89%</div> <div>10%</div> <div>..</div> </div>
1	B	291	<div> <div></div> <div>87%</div> <div>10%</div> <div>..</div> </div>
1	C	291	<div> <div></div> <div>87%</div> <div>10%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BR	A	401	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enterochelin uptake periplasmic binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	289	Total	C	N	O	S	0	7	0
			2196	1425	352	416	3			
1	B	288	Total	C	N	O	S	0	4	0
			2176	1416	343	414	3			
1	C	288	Total	C	N	O	S	0	4	0
			2159	1396	353	407	3			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	GLY	-	expression tag	UNP A0A0W8LI20
A	21	PRO	-	expression tag	UNP A0A0W8LI20
A	22	ALA	-	expression tag	UNP A0A0W8LI20
A	23	MET	-	expression tag	UNP A0A0W8LI20
A	227	LEU	HIS	engineered mutation	UNP A0A0W8LI20
B	20	GLY	-	expression tag	UNP A0A0W8LI20
B	21	PRO	-	expression tag	UNP A0A0W8LI20
B	22	ALA	-	expression tag	UNP A0A0W8LI20
B	23	MET	-	expression tag	UNP A0A0W8LI20
B	227	LEU	HIS	engineered mutation	UNP A0A0W8LI20
C	20	GLY	-	expression tag	UNP A0A0W8LI20
C	21	PRO	-	expression tag	UNP A0A0W8LI20
C	22	ALA	-	expression tag	UNP A0A0W8LI20
C	23	MET	-	expression tag	UNP A0A0W8LI20
C	227	LEU	HIS	engineered mutation	UNP A0A0W8LI20

- Molecule 2 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Br	0	0
			3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Br 1	0	0
2	C	1	Total 1	Br 1	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total 1	Na 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	123	Total 123	O 123	0	0
4	B	90	Total 90	O 90	0	0
4	C	88	Total 88	O 88	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enterochelin uptake periplasmic binding protein



- Molecule 1: Enterochelin uptake periplasmic binding protein



- Molecule 1: Enterochelin uptake periplasmic binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	56.92Å 62.56Å 67.79Å 82.21° 76.99° 76.18°	Depositor
Resolution (Å)	65.52 – 1.52 60.52 – 1.52	Depositor EDS
% Data completeness (in resolution range)	95.3 (65.52-1.52) 95.4 (60.52-1.52)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.185 , 0.221 0.194 , 0.228	Depositor DCC
R_{free} test set	6332 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6838	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.17	2/2246 (0.1%)	1.16	14/3038 (0.5%)
1	B	1.17	2/2220 (0.1%)	1.25	13/3005 (0.4%)
1	C	1.17	4/2194 (0.2%)	1.21	15/2971 (0.5%)
All	All	1.17	8/6660 (0.1%)	1.21	42/9014 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	160	GLU	CG-CD	6.80	1.62	1.51
1	B	243	TYR	CG-CD1	-6.54	1.30	1.39
1	B	291	SER	CB-OG	-6.32	1.34	1.42
1	C	206	PHE	CE2-CZ	5.81	1.48	1.37
1	C	93	SER	CB-OG	-5.33	1.35	1.42
1	A	160	GLU	CG-CD	5.29	1.59	1.51
1	C	152	SER	CA-CB	-5.21	1.45	1.52
1	A	142	PHE	CG-CD2	-5.12	1.31	1.38

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	301	MET	CG-SD-CE	-15.05	76.11	100.20
1	B	258	ARG	NE-CZ-NH2	-12.27	114.17	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	258	ARG	NE-CZ-NH1	10.58	125.59	120.30
1	B	23	MET	CA-CB-CG	8.77	128.22	113.30
1	B	60[A]	ILE	CG1-CB-CG2	-8.60	92.48	111.40
1	B	60[B]	ILE	CG1-CB-CG2	-8.60	92.48	111.40
1	C	65	ASP	CB-CG-OD2	-8.59	110.57	118.30
1	C	258	ARG	NE-CZ-NH1	8.45	124.53	120.30
1	A	301	MET	CG-SD-CE	-8.32	86.89	100.20
1	B	23	MET	CB-CG-SD	8.21	137.02	112.40
1	C	28	MET	CG-SD-CE	-7.53	88.15	100.20
1	C	157	TYR	CB-CG-CD2	7.53	125.52	121.00
1	C	258	ARG	NE-CZ-NH2	-7.49	116.56	120.30
1	A	81	LEU	CA-CB-CG	7.47	132.48	115.30
1	B	57	ASP	CB-CG-OD2	7.30	124.87	118.30
1	A	28	MET	CG-SD-CE	-7.24	88.61	100.20
1	C	81	LEU	CA-CB-CG	7.00	131.39	115.30
1	C	118	ARG	NE-CZ-NH2	6.80	123.70	120.30
1	C	33	ASP	CB-CG-OD1	6.73	124.36	118.30
1	B	24	LEU	CA-CB-CG	6.55	130.36	115.30
1	C	118	ARG	NE-CZ-NH1	-6.54	117.03	120.30
1	C	65	ASP	CB-CG-OD1	6.47	124.12	118.30
1	A	281	TYR	CB-CG-CD2	-6.39	117.17	121.00
1	A	124	ASP	CB-CG-OD1	6.25	123.92	118.30
1	A	249	ARG	NE-CZ-NH1	6.13	123.37	120.30
1	A	39	ASP	CB-CG-OD1	6.07	123.76	118.30
1	C	180	ILE	CG1-CB-CG2	5.95	124.48	111.40
1	A	65	ASP	CB-CG-OD1	5.62	123.36	118.30
1	B	62	ASP	CB-CG-OD1	5.59	123.33	118.30
1	A	283	ASP	CB-CG-OD2	-5.54	113.31	118.30
1	C	81	LEU	CB-CG-CD1	5.52	120.38	111.00
1	B	159	LEU	CB-CG-CD2	5.50	120.35	111.00
1	B	101	ASP	CB-CG-OD1	5.39	123.15	118.30
1	A	124	ASP	CB-CG-OD2	-5.38	113.46	118.30
1	A	62	ASP	CB-CG-OD1	5.34	123.11	118.30
1	A	101	ASP	CB-CG-OD1	5.30	123.07	118.30
1	A	249	ARG	NE-CZ-NH2	-5.29	117.66	120.30
1	C	211	ASP	CB-CG-OD1	5.21	122.99	118.30
1	C	24	LEU	CA-CB-CG	5.11	127.05	115.30
1	A	65	ASP	CB-CG-OD2	-5.08	113.73	118.30
1	B	124	ASP	CB-CG-OD1	5.08	122.87	118.30
1	C	165	GLU	CA-CB-CG	5.06	124.53	113.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	80	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2196	0	2216	26	0
1	B	2176	0	2158	36	0
1	C	2159	0	2124	28	0
2	A	3	0	0	2	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	C	1	0	0	0	0
4	A	123	0	0	10	1
4	B	90	0	0	13	0
4	C	88	0	0	9	1
All	All	6838	0	6498	90	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:MET:HA	4:B:578:HOH:O	1.13	1.26
1:A:192:THR:O	4:A:501:HOH:O	1.63	1.16
1:B:23:MET:CG	4:B:503:HOH:O	1.89	1.16
1:C:60[B]:ILE:HD12	1:C:114:ILE:HG21	1.31	1.11
1:C:119:GLN:OE1	4:C:501:HOH:O	1.67	1.11
1:C:258:ARG:NE	4:C:502:HOH:O	1.87	1.08
1:A:192:THR:C	4:A:501:HOH:O	1.93	1.05
1:C:119:GLN:CD	4:C:501:HOH:O	1.94	1.03
1:B:23:MET:HG2	4:B:503:HOH:O	1.54	0.98
1:C:119:GLN:NE2	4:C:501:HOH:O	1.98	0.97
1:A:159[B]:LEU:HD22	1:A:159[B]:LEU:N	1.85	0.91

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:60[B]:ILE:CD1	1:C:114:ILE:CG2	2.52	0.88
1:C:60[B]:ILE:HD12	1:C:114:ILE:CG2	2.04	0.87
1:B:60[B]:ILE:HG21	1:B:135:VAL:HG21	1.55	0.87
1:B:227:LEU:O	4:B:502:HOH:O	1.98	0.81
1:A:159[B]:LEU:N	1:A:159[B]:LEU:CD2	2.42	0.81
1:A:192:THR:OG1	4:A:501:HOH:O	1.99	0.80
1:B:23:MET:CB	4:B:503:HOH:O	2.23	0.79
1:A:103:GLU:CG	4:A:623:HOH:O	2.31	0.79
1:B:60[B]:ILE:CD1	1:B:114:ILE:HG21	2.13	0.79
1:A:26:ILE:H	1:A:148:ASN:HD21	1.29	0.79
1:C:26:ILE:H	1:C:148:ASN:HD21	1.32	0.78
1:A:147:GLU:CG	4:A:567:HOH:O	2.36	0.72
1:B:23:MET:N	4:B:503:HOH:O	2.23	0.72
1:B:55:ILE:HG23	1:B:60[B]:ILE:HD11	1.73	0.70
1:B:23:MET:HA	4:B:584:HOH:O	1.91	0.70
1:B:23:MET:CA	4:B:503:HOH:O	2.40	0.70
1:C:60[B]:ILE:CD1	1:C:114:ILE:HG22	2.21	0.70
1:C:277:LYS:CB	4:C:582:HOH:O	2.39	0.70
1:A:158:GLY:C	1:A:159[B]:LEU:HD22	2.12	0.69
1:A:70:ASN:H	1:A:70:ASN:HD22	1.42	0.68
1:B:60[B]:ILE:CD1	1:B:114:ILE:CG2	2.73	0.66
1:C:60[B]:ILE:CD1	1:C:114:ILE:HG21	2.08	0.66
1:B:89:LYS:CD	4:B:546:HOH:O	2.44	0.66
1:C:60[B]:ILE:HG21	1:C:135:VAL:HG21	1.77	0.65
1:B:26:ILE:H	1:B:148:ASN:HD21	1.44	0.64
1:C:258:ARG:CZ	4:C:502:HOH:O	2.36	0.62
1:A:157:TYR:O	1:A:159[B]:LEU:CD2	2.47	0.62
1:C:70:ASN:HD22	1:C:70:ASN:H	1.47	0.62
1:C:99:GLN:CG	4:C:581:HOH:O	2.48	0.62
1:B:194:SER:H	1:B:250:ASN:HD21	1.46	0.61
1:C:57:ASP:OD2	1:C:60[B]:ILE:HG23	2.00	0.60
1:C:258:ARG:NH2	4:C:502:HOH:O	2.36	0.59
1:C:250:ASN:HD22	1:C:255:ASN:HD22	1.51	0.59
1:B:250:ASN:HD22	1:B:255:ASN:HD22	1.51	0.59
1:B:57:ASP:OD2	1:B:60[B]:ILE:HG23	2.03	0.58
1:B:60[B]:ILE:HD13	1:B:114:ILE:CG2	2.34	0.58
1:B:210:HIS:HD2	1:B:215:ILE:O	1.87	0.57
1:B:70:ASN:H	1:B:70:ASN:HD22	1.52	0.57
1:B:60[B]:ILE:HD12	1:B:114:ILE:HG21	1.88	0.55
1:B:60[B]:ILE:HD13	1:B:114:ILE:HG21	1.86	0.55
1:B:23:MET:CA	4:B:578:HOH:O	1.95	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:ILE:N	1:B:148:ASN:HD21	2.05	0.54
1:B:194:SER:N	1:B:250:ASN:HD21	2.04	0.54
1:A:234:GLU:CG	4:A:575:HOH:O	2.56	0.53
1:C:60[B]:ILE:HD11	1:C:114:ILE:CG2	2.36	0.52
1:B:23:MET:N	4:B:505:HOH:O	2.42	0.51
1:C:210:HIS:HD2	1:C:215:ILE:O	1.94	0.51
1:B:134:PHE:CE2	1:B:136:GLY:HA3	2.46	0.51
1:C:220:GLU:CB	4:C:584:HOH:O	2.58	0.50
1:A:192:THR:CB	4:A:501:HOH:O	2.54	0.49
1:C:194:SER:N	1:C:250:ASN:HD21	2.11	0.47
1:C:60[B]:ILE:HD11	1:C:114:ILE:HG22	1.94	0.47
1:A:210:HIS:HE1	4:A:530:HOH:O	1.97	0.46
1:C:194:SER:H	1:C:250:ASN:HD21	1.63	0.46
1:A:39:ASP:OD2	2:A:401:BR:BR	2.89	0.46
1:B:23:MET:HB2	1:B:138:ASP:OD2	2.15	0.46
1:B:227:LEU:CB	4:B:504:HOH:O	2.64	0.46
1:B:255:ASN:O	1:B:258:ARG:NH2	2.50	0.45
1:B:55:ILE:O	1:B:77:PRO:HD3	2.17	0.45
1:A:37:VAL:HG21	1:A:46:ILE:HD12	1.98	0.45
1:A:193:ASN:HB3	4:A:521:HOH:O	2.15	0.45
1:C:60[B]:ILE:HD13	1:C:135:VAL:HG21	1.99	0.45
1:A:192:THR:CA	4:A:501:HOH:O	2.52	0.44
1:A:26:ILE:N	1:A:148:ASN:HD21	2.06	0.44
1:A:70:ASN:H	1:A:70:ASN:ND2	2.12	0.43
1:B:60[B]:ILE:CG2	1:B:135:VAL:HG21	2.37	0.43
1:A:157:TYR:O	1:A:159[B]:LEU:HD21	2.19	0.42
1:A:248:ASP:OD1	1:A:250:ASN:HB2	2.18	0.42
1:A:37:VAL:HG12	2:A:401:BR:BR	2.75	0.42
1:A:159[B]:LEU:N	1:A:159[B]:LEU:HD23	2.33	0.42
1:B:60[A]:ILE:HG23	1:B:60[A]:ILE:HD12	1.52	0.42
1:A:210:HIS:HD2	1:A:215:ILE:O	2.02	0.42
1:B:80:ASN:OD1	1:B:97:VAL:N	2.46	0.42
1:C:142:PHE:CD1	1:C:142:PHE:C	2.92	0.41
1:B:70:ASN:H	1:B:70:ASN:ND2	2.17	0.41
1:A:180:ILE:C	1:A:180:ILE:HD12	2.42	0.40
1:C:57:ASP:HB3	1:C:60[B]:ILE:CD1	2.51	0.40
1:B:210:HIS:HE1	4:B:536:HOH:O	2.04	0.40
1:C:159[A]:LEU:HD12	1:C:159[A]:LEU:HA	1.92	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:615:HOH:O	4:C:555:HOH:O[1_455]	1.38	0.82

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/291 (101%)	284 (97%)	10 (3%)	0	100	100
1	B	290/291 (100%)	282 (97%)	7 (2%)	1 (0%)	41	18
1	C	290/291 (100%)	279 (96%)	11 (4%)	0	100	100
All	All	874/873 (100%)	845 (97%)	28 (3%)	1 (0%)	51	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	98	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/252 (90%)	225 (99%)	3 (1%)	69	43
1	B	218/252 (86%)	210 (96%)	8 (4%)	34	8
1	C	215/252 (85%)	206 (96%)	9 (4%)	30	5
All	All	661/756 (87%)	641 (97%)	20 (3%)	47	12

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	159[A]	LEU
1	A	159[B]	LEU
1	B	23	MET
1	B	24	LEU
1	B	36	LEU
1	B	51	SER
1	B	146	PHE
1	B	159	LEU
1	B	195[A]	ASN
1	B	195[B]	ASN
1	C	60[A]	ILE
1	C	60[B]	ILE
1	C	70	ASN
1	C	81	LEU
1	C	159[A]	LEU
1	C	159[B]	LEU
1	C	180	ILE
1	C	256	LYS
1	C	271	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	90	ASN
1	A	148	ASN
1	A	172	ASN
1	A	210	HIS
1	A	250	ASN
1	B	70	ASN
1	B	148	ASN
1	B	210	HIS
1	B	250	ASN
1	C	49	ASN
1	C	70	ASN
1	C	90	ASN
1	C	98	GLN
1	C	148	ASN
1	C	210	HIS
1	C	216	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	250	ASN
1	C	307	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/291 (99%)	-0.50	2 (0%) 87 90	16, 25, 39, 55	3 (1%)
1	B	288/291 (98%)	-0.49	3 (1%) 82 85	17, 26, 40, 71	4 (1%)
1	C	288/291 (98%)	-0.41	1 (0%) 94 95	16, 27, 41, 59	3 (1%)
All	All	865/873 (99%)	-0.47	6 (0%) 87 90	16, 26, 41, 71	10 (1%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	222	ILE	3.8
1	A	222	ILE	3.7
1	B	224	VAL	2.7
1	A	97	VAL	2.6
1	C	224	VAL	2.3
1	B	225	GLY	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BR	B	401	1/1	0.73	0.10	133,133,133,133	0
2	BR	A	401	1/1	0.94	0.14	86,86,86,86	0
3	NA	C	402	1/1	0.98	0.09	23,23,23,23	0
2	BR	C	401	1/1	0.99	0.02	21,21,21,21	1
2	BR	A	402	1/1	0.99	0.03	25,25,25,25	1
2	BR	A	403	1/1	1.00	0.03	25,25,25,25	1

6.5 Other polymers [i](#)

There are no such residues in this entry.