



# Full wwPDB X-ray Structure Validation Report ⓘ

May 28, 2020 – 07:59 pm BST

PDB ID : 1MA8  
Title : A-DNA decamer GCGTA(UMS)ACGC with incorporated 2'-methylseleno-uridine  
Authors : Teplova, M.; Wilds, C.J.; Wawrzak, Z.; Tereshko, V.; Du, Q.; Carrasco, N.; Huang, Z.; Egli, M.  
Deposited on : 2002-08-01  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 769 atoms, of which 222 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*GP\*CP\*GP\*TP\*AP\*UMSP\*AP\*CP\*GP\*C)-3'.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P	Se			
1	A	10	314	97	111	38	58	9	1	0	0	0
1	B	10	314	97	111	38	58	9	1	0	0	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	72	Total	O	0	0
			72	72		
2	B	69	Total	O	0	0
			69	69		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 5'-D(\*GP\*CP\*GP\*TP\*AP\*UMSP\*AP\*CP\*GP\*C)-3'

Chain A:  100%

G1	C2	G3	T4	A5	U6	A7	C8	G9	C10
----	----	----	----	----	----	----	----	----	-----

- Molecule 1: 5'-D(\*GP\*CP\*GP\*TP\*AP\*UMSP\*AP\*CP\*GP\*C)-3'

Chain B:  10% 90%

G111	C112	G113	T114	A115	U116	A117	C118	G119	C120
------	------	------	------	------	------	------	------	------	------

## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	24.55Å 43.97Å 45.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.30	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.30)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
Refinement program	SHELXL-97, CNS	Depositor
R, $R_{free}$	0.149 , 0.173	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	769	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:  
UMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	2.32	10/204 (4.9%)	2.37	15/313 (4.8%)
1	B	2.71	15/204 (7.4%)	2.63	22/313 (7.0%)
All	All	2.52	25/408 (6.1%)	2.51	37/626 (5.9%)

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	119	DG	C2'-C1'	-9.97	1.42	1.52
1	B	120	DC	C2'-C1'	-8.34	1.44	1.52
1	A	4	DT	C2'-C1'	-8.24	1.44	1.52
1	B	115	DA	C2'-C1'	-8.02	1.44	1.52
1	A	2	DC	N1-C6	-7.66	1.32	1.37
1	A	3	DG	C2'-C1'	-7.39	1.44	1.52
1	B	119	DG	O4'-C1'	7.30	1.51	1.42
1	B	111	DG	C2'-C1'	-7.29	1.45	1.52
1	B	120	DC	O4'-C1'	7.18	1.50	1.42
1	A	2	DC	C2'-C1'	-7.00	1.45	1.52
1	B	112	DC	C4'-O4'	6.78	1.51	1.45
1	A	1	DG	C8-N7	6.72	1.34	1.30
1	A	8	DC	N1-C6	-6.50	1.33	1.37
1	B	112	DC	O4'-C1'	6.36	1.49	1.42
1	B	112	DC	C3'-O3'	6.24	1.52	1.44
1	A	5	DA	N3-C4	-6.21	1.31	1.34
1	A	9	DG	C2'-C1'	-6.20	1.46	1.52
1	B	115	DA	O4'-C1'	6.16	1.49	1.42
1	B	112	DC	C2'-C1'	-6.16	1.46	1.52
1	B	113	DG	C4'-O4'	5.91	1.50	1.45
1	B	119	DG	N9-C8	-5.87	1.33	1.37
1	A	1	DG	C2'-C1'	-5.50	1.46	1.52
1	B	112	DC	C4'-C3'	-5.30	1.47	1.52
1	A	3	DG	O4'-C1'	5.21	1.48	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	119	DG	C3'-C2'	-5.17	1.46	1.52

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	DG	O4'-C1'-N9	-9.31	101.48	108.00
1	B	120	DC	O4'-C1'-N1	-8.39	102.12	108.00
1	B	115	DA	O4'-C1'-N9	-8.13	102.31	108.00
1	A	9	DG	O4'-C1'-N9	-7.68	102.62	108.00
1	B	112	DC	C1'-O4'-C4'	-7.62	102.48	110.10
1	A	1	DG	O4'-C1'-N9	-7.57	102.70	108.00
1	B	113	DG	N7-C8-N9	7.47	116.84	113.10
1	B	119	DG	O4'-C1'-N9	-7.39	102.82	108.00
1	A	2	DC	C5-C6-N1	7.30	124.65	121.00
1	B	111	DG	O4'-C1'-N9	-7.28	102.90	108.00
1	A	7	DA	O4'-C4'-C3'	-6.99	101.70	104.50
1	A	10	DC	N3-C4-N4	-6.80	113.24	118.00
1	B	113	DG	C5-C6-O6	6.56	132.53	128.60
1	B	113	DG	C5-N7-C8	-6.49	101.06	104.30
1	A	8	DC	O4'-C4'-C3'	-6.43	101.93	104.50
1	A	9	DG	N1-C6-O6	6.39	123.74	119.90
1	B	113	DG	N1-C6-O6	-6.38	116.07	119.90
1	B	114	DT	C4-C5-C7	6.34	122.81	119.00
1	B	117	DA	P-O5'-C5'	-6.31	110.80	120.90
1	A	2	DC	C4-C5-C6	-6.23	114.28	117.40
1	A	4	DT	O4'-C1'-N1	-6.22	103.65	108.00
1	B	114	DT	C5-C6-N1	6.19	127.41	123.70
1	B	115	DA	C8-N9-C4	6.13	108.25	105.80
1	A	9	DG	C8-N9-C1'	5.99	134.78	127.00
1	A	5	DA	C2-N3-C4	5.69	113.44	110.60
1	A	9	DG	N7-C8-N9	5.64	115.92	113.10
1	B	117	DA	C4'-C3'-C2'	-5.43	98.21	103.10
1	B	112	DC	C4-C5-C6	5.43	120.11	117.40
1	B	118	DC	C6-N1-C2	-5.37	118.15	120.30
1	A	9	DG	C5-C6-O6	-5.19	125.49	128.60
1	B	119	DG	N1-C6-O6	-5.16	116.80	119.90
1	B	119	DG	N7-C8-N9	5.08	115.64	113.10
1	A	9	DG	C8-N9-C4	-5.04	104.38	106.40
1	B	111	DG	C1'-O4'-C4'	-5.03	105.07	110.10
1	B	113	DG	N3-C4-C5	-5.03	126.09	128.60
1	B	118	DC	C5-C6-N1	5.01	123.50	121.00
1	B	115	DA	N7-C8-N9	-5.00	111.30	113.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	203	111	114	3	0
1	B	203	111	114	0	0
2	A	72	0	0	3	2
2	B	69	0	0	0	1
All	All	547	222	228	3	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:UMS:SE2'	2:A:837:HOH:O	2.63	0.67
1:A:6:UMS:H1'	2:A:837:HOH:O	2.12	0.49
1:A:6:UMS:C1'	2:A:837:HOH:O	2.67	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:846:HOH:O	2:B:842:HOH:O[1_455]	1.95	0.25
2:A:752:HOH:O	2:A:808:HOH:O[3_655]	2.02	0.18

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	UMS	A	6	1	13,22,23	2.41	6 (46%)	11,31,34	1.98	4 (36%)
1	UMS	B	116	1	13,22,23	1.77	4 (30%)	11,31,34	2.08	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	UMS	A	6	1	-	0/5/27/28	0/2/2/2
1	UMS	B	116	1	-	0/5/27/28	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	UMS	C2'-C3'	-4.68	1.47	1.53
1	B	116	UMS	O5'-C5'	3.91	1.54	1.44
1	A	6	UMS	C4-N3	3.65	1.39	1.33
1	A	6	UMS	O5'-C5'	3.36	1.53	1.44
1	A	6	UMS	C5'-C4'	-3.04	1.42	1.51
1	B	116	UMS	O4'-C4'	2.90	1.51	1.45

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	UMS	C6-N1	2.84	1.39	1.35
1	A	6	UMS	O4'-C4'	2.46	1.50	1.45
1	B	116	UMS	C5'-C4'	-2.44	1.44	1.51
1	B	116	UMS	C6-N1	2.24	1.38	1.35

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	116	UMS	C5-C4-N3	-4.85	112.64	123.31
1	B	116	UMS	O4'-C4'-C3'	-4.08	97.04	105.11
1	A	6	UMS	O3'-C3'-C2'	3.67	116.47	110.64
1	A	6	UMS	O4'-C4'-C3'	-3.65	97.90	105.11
1	A	6	UMS	C5-C4-N3	-3.13	116.42	123.31
1	A	6	UMS	O3'-C3'-C4'	-2.14	104.87	111.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	6	UMS	3	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.