



wwPDB X-ray Structure Validation Summary Report i

Sep 11, 2023 – 06:49 PM EDT

PDB ID : 4MHI
Title : Crystal structure of a H5N1 influenza virus hemagglutinin from A/goose/Guangdong/1/96
Authors : Zhu, X.; Wilson, I.A.
Deposited on : 2013-08-29
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	K	334	4%	72%	23%	..
1	M	334	8%	75%	20%	..
1	O	334	5%	76%	20%	..
1	Q	334	%	78%	17%	..
2	B	182	2%	80%	15%	5%
2	D	182	2%	77%	18%	5%
2	F	182	10%	78%	17%	5%
2	H	182	8%	77%	17%	. 5%
2	J	182	%	86%	9%	5%
2	L	182	6%	76%	18%	. 5%
2	N	182	3%	83%	12%	. 5%
2	P	182	3%	80%	15%	. 5%
2	R	182	%	82%	13%	5%
3	S	2		100%		
3	T	2		100%		
3	U	2		100%		
3	V	2		50%	50%	
3	W	2		50%	50%	
3	X	2		50%	50%	
3	Z	2		100%		
4	Y	3		100%		

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 36134 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0
1	C	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0
1	E	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0
1	G	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0
1	I	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0
1	K	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0
1	M	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0
1	O	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0
1	Q	323	Total 2551	C 1609	N 442	O 485	S 15	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ALA	-	expression tag	UNP Q9Q0U6
A	8	ASP	-	expression tag	UNP Q9Q0U6
A	9	PRO	-	expression tag	UNP Q9Q0U6
A	10	GLY	-	expression tag	UNP Q9Q0U6
C	7	ALA	-	expression tag	UNP Q9Q0U6
C	8	ASP	-	expression tag	UNP Q9Q0U6
C	9	PRO	-	expression tag	UNP Q9Q0U6
C	10	GLY	-	expression tag	UNP Q9Q0U6
E	7	ALA	-	expression tag	UNP Q9Q0U6
E	8	ASP	-	expression tag	UNP Q9Q0U6
E	9	PRO	-	expression tag	UNP Q9Q0U6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	10	GLY	-	expression tag	UNP Q9Q0U6
G	7	ALA	-	expression tag	UNP Q9Q0U6
G	8	ASP	-	expression tag	UNP Q9Q0U6
G	9	PRO	-	expression tag	UNP Q9Q0U6
G	10	GLY	-	expression tag	UNP Q9Q0U6
I	7	ALA	-	expression tag	UNP Q9Q0U6
I	8	ASP	-	expression tag	UNP Q9Q0U6
I	9	PRO	-	expression tag	UNP Q9Q0U6
I	10	GLY	-	expression tag	UNP Q9Q0U6
K	7	ALA	-	expression tag	UNP Q9Q0U6
K	8	ASP	-	expression tag	UNP Q9Q0U6
K	9	PRO	-	expression tag	UNP Q9Q0U6
K	10	GLY	-	expression tag	UNP Q9Q0U6
M	7	ALA	-	expression tag	UNP Q9Q0U6
M	8	ASP	-	expression tag	UNP Q9Q0U6
M	9	PRO	-	expression tag	UNP Q9Q0U6
M	10	GLY	-	expression tag	UNP Q9Q0U6
O	7	ALA	-	expression tag	UNP Q9Q0U6
O	8	ASP	-	expression tag	UNP Q9Q0U6
O	9	PRO	-	expression tag	UNP Q9Q0U6
O	10	GLY	-	expression tag	UNP Q9Q0U6
Q	7	ALA	-	expression tag	UNP Q9Q0U6
Q	8	ASP	-	expression tag	UNP Q9Q0U6
Q	9	PRO	-	expression tag	UNP Q9Q0U6
Q	10	GLY	-	expression tag	UNP Q9Q0U6

- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	173	Total	C	N	O	S			
			1403	872	242	281	8	0	0	0
2	D	173	Total	C	N	O	S			
			1403	872	242	281	8	0	0	0
2	F	173	Total	C	N	O	S			
			1403	872	242	281	8	0	0	0
2	H	173	Total	C	N	O	S			
			1403	872	242	281	8	0	0	0
2	J	173	Total	C	N	O	S			
			1403	872	242	281	8	0	0	0
2	L	173	Total	C	N	O	S			
			1403	872	242	281	8	0	0	0
2	N	173	Total	C	N	O	S			
			1403	872	242	281	8	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	173	Total	C	N	O	S	0	0	0
			1403	872	242	281	8			

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	176	SER	-	expression tag	UNP Q9Q0U6
B	177	GLY	-	expression tag	UNP Q9Q0U6
B	178	ARG	-	expression tag	UNP Q9Q0U6
B	179	LEU	-	expression tag	UNP Q9Q0U6
B	180	VAL	-	expression tag	UNP Q9Q0U6
B	181	PRO	-	expression tag	UNP Q9Q0U6
B	182	ARG	-	expression tag	UNP Q9Q0U6
D	176	SER	-	expression tag	UNP Q9Q0U6
D	177	GLY	-	expression tag	UNP Q9Q0U6
D	178	ARG	-	expression tag	UNP Q9Q0U6
D	179	LEU	-	expression tag	UNP Q9Q0U6
D	180	VAL	-	expression tag	UNP Q9Q0U6
D	181	PRO	-	expression tag	UNP Q9Q0U6
D	182	ARG	-	expression tag	UNP Q9Q0U6
F	176	SER	-	expression tag	UNP Q9Q0U6
F	177	GLY	-	expression tag	UNP Q9Q0U6
F	178	ARG	-	expression tag	UNP Q9Q0U6
F	179	LEU	-	expression tag	UNP Q9Q0U6
F	180	VAL	-	expression tag	UNP Q9Q0U6
F	181	PRO	-	expression tag	UNP Q9Q0U6
F	182	ARG	-	expression tag	UNP Q9Q0U6
H	176	SER	-	expression tag	UNP Q9Q0U6
H	177	GLY	-	expression tag	UNP Q9Q0U6
H	178	ARG	-	expression tag	UNP Q9Q0U6
H	179	LEU	-	expression tag	UNP Q9Q0U6
H	180	VAL	-	expression tag	UNP Q9Q0U6
H	181	PRO	-	expression tag	UNP Q9Q0U6
H	182	ARG	-	expression tag	UNP Q9Q0U6
J	176	SER	-	expression tag	UNP Q9Q0U6
J	177	GLY	-	expression tag	UNP Q9Q0U6
J	178	ARG	-	expression tag	UNP Q9Q0U6
J	179	LEU	-	expression tag	UNP Q9Q0U6
J	180	VAL	-	expression tag	UNP Q9Q0U6
J	181	PRO	-	expression tag	UNP Q9Q0U6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	182	ARG	-	expression tag	UNP Q9Q0U6
L	176	SER	-	expression tag	UNP Q9Q0U6
L	177	GLY	-	expression tag	UNP Q9Q0U6
L	178	ARG	-	expression tag	UNP Q9Q0U6
L	179	LEU	-	expression tag	UNP Q9Q0U6
L	180	VAL	-	expression tag	UNP Q9Q0U6
L	181	PRO	-	expression tag	UNP Q9Q0U6
L	182	ARG	-	expression tag	UNP Q9Q0U6
N	176	SER	-	expression tag	UNP Q9Q0U6
N	177	GLY	-	expression tag	UNP Q9Q0U6
N	178	ARG	-	expression tag	UNP Q9Q0U6
N	179	LEU	-	expression tag	UNP Q9Q0U6
N	180	VAL	-	expression tag	UNP Q9Q0U6
N	181	PRO	-	expression tag	UNP Q9Q0U6
N	182	ARG	-	expression tag	UNP Q9Q0U6
P	176	SER	-	expression tag	UNP Q9Q0U6
P	177	GLY	-	expression tag	UNP Q9Q0U6
P	178	ARG	-	expression tag	UNP Q9Q0U6
P	179	LEU	-	expression tag	UNP Q9Q0U6
P	180	VAL	-	expression tag	UNP Q9Q0U6
P	181	PRO	-	expression tag	UNP Q9Q0U6
P	182	ARG	-	expression tag	UNP Q9Q0U6
R	176	SER	-	expression tag	UNP Q9Q0U6
R	177	GLY	-	expression tag	UNP Q9Q0U6
R	178	ARG	-	expression tag	UNP Q9Q0U6
R	179	LEU	-	expression tag	UNP Q9Q0U6
R	180	VAL	-	expression tag	UNP Q9Q0U6
R	181	PRO	-	expression tag	UNP Q9Q0U6
R	182	ARG	-	expression tag	UNP Q9Q0U6

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	S	2	Total C N O				0	0	0
			28 16 2 10						
3	T	2	Total C N O				0	0	0
			28 16 2 10						

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	U	2	Total C N O 28 16 2 10	0	0	0
3	V	2	Total C N O 28 16 2 10	0	0	0
3	W	2	Total C N O 28 16 2 10	0	0	0
3	X	2	Total C N O 28 16 2 10	0	0	0
3	Z	2	Total C N O 28 16 2 10	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	Y	3	Total C N O 39 22 2 15	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 14 8 1 5	0	0
5	C	1	Total C N O 14 8 1 5	0	0
5	E	1	Total C N O 14 8 1 5	0	0
5	E	1	Total C N O 14 8 1 5	0	0
5	G	1	Total C N O 14 8 1 5	0	0
5	G	1	Total C N O 14 8 1 5	0	0
5	I	1	Total C N O 14 8 1 5	0	0
5	K	1	Total C N O 14 8 1 5	0	0
5	K	1	Total C N O 14 8 1 5	0	0
5	M	1	Total C N O 14 8 1 5	0	0
5	M	1	Total C N O 14 8 1 5	0	0
5	O	1	Total C N O 14 8 1 5	0	0
5	O	1	Total C N O 14 8 1 5	0	0
5	Q	1	Total C N O 14 8 1 5	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	7	Total O 7 7	0	0
6	B	5	Total O 5 5	0	0
6	C	6	Total O 6 6	0	0
6	D	8	Total O 8 8	0	0
6	E	6	Total O 6 6	0	0
6	F	2	Total O 2 2	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	G	14	Total O 14 14	0	0
6	H	4	Total O 4 4	0	0
6	I	9	Total O 9 9	0	0
6	J	12	Total O 12 12	0	0
6	K	10	Total O 10 10	0	0
6	L	2	Total O 2 2	0	0
6	M	5	Total O 5 5	0	0
6	N	1	Total O 1 1	0	0
6	O	8	Total O 8 8	0	0
6	P	2	Total O 2 2	0	0
6	Q	8	Total O 8 8	0	0
6	R	8	Total O 8 8	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin HA1 chain



- Molecule 1: Hemagglutinin HA1 chain





- Molecule 1: Hemagglutinin HA1 chain







- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 2: Hemagglutinin HA2 chain




- Molecule 2: Hemagglutinin HA2 chain




- Molecule 2: Hemagglutinin HA2 chain




- Molecule 2: Hemagglutinin HA2 chain





- Molecule 2: Hemagglutinin HA2 chain





- Molecule 2: Hemagglutinin HA2 chain





- Molecule 2: Hemagglutinin HA2 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.48Å 225.73Å 211.63Å 90.00° 98.97° 90.00°	Depositor
Resolution (Å)	49.66 – 2.60 49.66 – 2.60	Depositor EDS
% Data completeness (in resolution range)	94.8 (49.66-2.60) 95.1 (49.66-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ ¹	1.17 (at 2.58Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R , R_{free}	0.189 , 0.243 0.199 , 0.248	Depositor DCC
R_{free} test set	9880 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	65.9	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 63.0	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	36134	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.26 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7239e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2616	0.67	1/3557 (0.0%)
1	C	0.44	0/2616	0.66	2/3557 (0.1%)
1	E	0.44	0/2616	0.65	2/3557 (0.1%)
1	G	0.46	0/2616	0.66	2/3557 (0.1%)
1	I	0.49	0/2616	0.70	1/3557 (0.0%)
1	K	0.44	0/2616	0.64	1/3557 (0.0%)
1	M	0.40	0/2616	0.62	2/3557 (0.1%)
1	O	0.42	0/2616	0.64	1/3557 (0.0%)
1	Q	0.46	0/2616	0.67	1/3557 (0.0%)
2	B	0.49	0/1430	0.65	0/1924
2	D	0.46	0/1430	0.58	0/1924
2	F	0.47	0/1430	0.60	0/1924
2	H	0.47	0/1430	0.60	0/1924
2	J	0.47	0/1430	0.61	0/1924
2	L	0.45	0/1430	0.55	0/1924
2	N	0.43	0/1430	0.59	0/1924
2	P	0.46	0/1430	0.60	0/1924
2	R	0.49	0/1430	0.62	0/1924
All	All	0.45	0/36414	0.64	13/49329 (0.0%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	I	9	PRO	CA-N-CD	-8.94	98.98	111.50
1	C	9	PRO	CA-N-CD	-8.81	99.17	111.50
1	O	9	PRO	CA-N-CD	-8.72	99.29	111.50
1	G	9	PRO	CA-N-CD	-8.66	99.38	111.50
1	K	9	PRO	CA-N-CD	-8.51	99.59	111.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2551	0	2476	45	0
1	C	2551	0	2476	55	0
1	E	2551	0	2474	54	0
1	G	2551	0	2476	54	0
1	I	2551	0	2476	49	0
1	K	2551	0	2475	58	0
1	M	2551	0	2475	60	0
1	O	2551	0	2476	48	0
1	Q	2551	0	2476	43	0
2	B	1403	0	1302	22	0
2	D	1403	0	1302	24	0
2	F	1403	0	1302	24	0
2	H	1403	0	1302	31	0
2	J	1403	0	1302	13	0
2	L	1403	0	1302	24	0
2	N	1403	0	1302	21	0
2	P	1403	0	1302	22	0
2	R	1403	0	1301	23	0
3	S	28	0	25	0	0
3	T	28	0	25	0	0
3	U	28	0	25	3	0
3	V	28	0	25	1	0
3	W	28	0	25	1	0
3	X	28	0	25	1	0
3	Z	28	0	25	0	0
4	Y	39	0	34	0	0
5	A	14	0	13	0	0
5	C	14	0	13	0	0
5	E	28	0	26	0	0
5	G	28	0	26	0	0
5	I	14	0	13	0	0
5	K	28	0	26	3	0
5	M	28	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	O	28	0	26	0	0
5	Q	14	0	13	0	0
6	A	7	0	0	1	0
6	B	5	0	0	0	0
6	C	6	0	0	0	0
6	D	8	0	0	0	0
6	E	6	0	0	0	0
6	F	2	0	0	0	0
6	G	14	0	0	0	0
6	H	4	0	0	0	0
6	I	9	0	0	0	0
6	J	12	0	0	0	0
6	K	10	0	0	3	0
6	L	2	0	0	0	0
6	M	5	0	0	0	0
6	N	1	0	0	0	0
6	O	8	0	0	1	0
6	P	2	0	0	0	0
6	Q	8	0	0	0	0
6	R	8	0	0	0	0
All	All	36134	0	34388	578	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 578 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:9:PRO:HD2	1:M:10:GLY:H	1.17	1.07
1:E:9:PRO:HD2	1:E:10:GLY:H	1.20	1.05
1:A:9:PRO:HD2	1:A:10:GLY:H	1.23	0.99
1:I:170:ASN:CG	1:I:176:LEU:HD23	1.85	0.97
1:I:170:ASN:ND2	1:I:176:LEU:HD23	1.81	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/334 (96%)	312 (97%)	9 (3%)	0	100 100
1	C	321/334 (96%)	309 (96%)	12 (4%)	0	100 100
1	E	321/334 (96%)	311 (97%)	10 (3%)	0	100 100
1	G	321/334 (96%)	311 (97%)	10 (3%)	0	100 100
1	I	321/334 (96%)	310 (97%)	11 (3%)	0	100 100
1	K	321/334 (96%)	311 (97%)	9 (3%)	1 (0%)	41 64
1	M	321/334 (96%)	309 (96%)	12 (4%)	0	100 100
1	O	321/334 (96%)	314 (98%)	7 (2%)	0	100 100
1	Q	321/334 (96%)	311 (97%)	10 (3%)	0	100 100
2	B	171/182 (94%)	167 (98%)	4 (2%)	0	100 100
2	D	171/182 (94%)	167 (98%)	4 (2%)	0	100 100
2	F	171/182 (94%)	166 (97%)	5 (3%)	0	100 100
2	H	171/182 (94%)	168 (98%)	3 (2%)	0	100 100
2	J	171/182 (94%)	168 (98%)	3 (2%)	0	100 100
2	L	171/182 (94%)	166 (97%)	5 (3%)	0	100 100
2	N	171/182 (94%)	167 (98%)	3 (2%)	1 (1%)	25 47
2	P	171/182 (94%)	167 (98%)	4 (2%)	0	100 100
2	R	171/182 (94%)	167 (98%)	4 (2%)	0	100 100
All	All	4428/4644 (95%)	4301 (97%)	125 (3%)	2 (0%)	100 100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	K	142	HIS
2	N	12	GLY

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	287/297 (97%)	277 (96%)	10 (4%)	36 62
1	C	287/297 (97%)	273 (95%)	14 (5%)	25 48
1	E	287/297 (97%)	272 (95%)	15 (5%)	23 46
1	G	287/297 (97%)	274 (96%)	13 (4%)	27 52
1	I	287/297 (97%)	276 (96%)	11 (4%)	33 59
1	K	287/297 (97%)	277 (96%)	10 (4%)	36 62
1	M	287/297 (97%)	277 (96%)	10 (4%)	36 62
1	O	287/297 (97%)	276 (96%)	11 (4%)	33 59
1	Q	287/297 (97%)	278 (97%)	9 (3%)	40 66
2	B	148/155 (96%)	147 (99%)	1 (1%)	84 94
2	D	148/155 (96%)	147 (99%)	1 (1%)	84 94
2	F	148/155 (96%)	147 (99%)	1 (1%)	84 94
2	H	148/155 (96%)	146 (99%)	2 (1%)	67 85
2	J	148/155 (96%)	146 (99%)	2 (1%)	67 85
2	L	148/155 (96%)	141 (95%)	7 (5%)	26 50
2	N	148/155 (96%)	147 (99%)	1 (1%)	84 94
2	P	148/155 (96%)	145 (98%)	3 (2%)	55 78
2	R	148/155 (96%)	147 (99%)	1 (1%)	84 94
All	All	3915/4068 (96%)	3793 (97%)	122 (3%)	40 66

5 of 122 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	23	THR
2	P	73	LEU
1	K	26	VAL
2	P	18	VAL
1	Q	151	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	197	ASN
1	M	211	GLN
1	Q	38	HIS
2	N	83	GLN
1	E	211	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	S	1	1,3	14,14,15	0.49	0	17,19,21	1.16	1 (5%)
3	NAG	S	2	3	14,14,15	0.52	0	17,19,21	0.98	1 (5%)
3	NAG	T	1	1,3	14,14,15	0.47	0	17,19,21	1.07	1 (5%)
3	NAG	T	2	3	14,14,15	0.42	0	17,19,21	0.83	1 (5%)
3	NAG	U	1	1,3	14,14,15	0.55	0	17,19,21	1.84	4 (23%)
3	NAG	U	2	3	14,14,15	0.52	0	17,19,21	1.38	2 (11%)
3	NAG	V	1	1,3	14,14,15	0.59	0	17,19,21	1.66	4 (23%)
3	NAG	V	2	3	14,14,15	0.54	0	17,19,21	1.00	1 (5%)
3	NAG	W	1	1,3	14,14,15	0.51	0	17,19,21	1.61	3 (17%)
3	NAG	W	2	3	14,14,15	0.54	0	17,19,21	0.70	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	X	1	1,3	14,14,15	0.54	0	17,19,21	1.61	3 (17%)
3	NAG	X	2	3	14,14,15	0.57	0	17,19,21	1.33	1 (5%)
4	NAG	Y	1	4,1	14,14,15	0.46	0	17,19,21	1.18	1 (5%)
4	NAG	Y	2	4	14,14,15	0.49	0	17,19,21	0.96	1 (5%)
4	MAN	Y	3	4	11,11,12	0.52	0	15,15,17	1.66	3 (20%)
3	NAG	Z	1	2,3	14,14,15	0.62	0	17,19,21	1.58	3 (17%)
3	NAG	Z	2	3	14,14,15	0.53	0	17,19,21	1.70	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	S	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	S	2	3	-	4/6/23/26	0/1/1/1
3	NAG	T	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	T	2	3	-	4/6/23/26	0/1/1/1
3	NAG	U	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	U	2	3	-	3/6/23/26	0/1/1/1
3	NAG	V	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	V	2	3	-	0/6/23/26	0/1/1/1
3	NAG	W	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	W	2	3	-	1/6/23/26	0/1/1/1
3	NAG	X	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	X	2	3	-	1/6/23/26	0/1/1/1
4	NAG	Y	1	4,1	-	4/6/23/26	0/1/1/1
4	NAG	Y	2	4	-	0/6/23/26	0/1/1/1
4	MAN	Y	3	4	-	2/2/19/22	0/1/1/1
3	NAG	Z	1	2,3	-	3/6/23/26	0/1/1/1
3	NAG	Z	2	3	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	V	1	NAG	C1-O5-C5	4.62	118.44	112.19
3	Z	2	NAG	C1-O5-C5	4.61	118.44	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	1	NAG	C1-O5-C5	4.43	118.20	112.19
3	U	2	NAG	C1-O5-C5	4.18	117.85	112.19
3	U	1	NAG	C3-C4-C5	4.06	117.48	110.24

There are no chirality outliers.

5 of 33 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	T	2	NAG	O7-C7-N2-C2
3	U	2	NAG	C8-C7-N2-C2
3	U	2	NAG	O7-C7-N2-C2
3	T	2	NAG	C8-C7-N2-C2
3	S	1	NAG	C8-C7-N2-C2


There are no ring outliers.


5 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	U	1	NAG	3	0
3	U	2	NAG	3	0
3	X	1	NAG	1	0
3	W	1	NAG	1	0
3	V	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




















5.6 Ligand geometry (i)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	E	2004	1	14,14,15	0.40	0	17,19,21	0.77	0
5	NAG	M	2003	1	14,14,15	0.45	0	17,19,21	0.82	0
5	NAG	O	2001	1	14,14,15	0.47	0	17,19,21	1.49	2 (11%)
5	NAG	M	2004	1	14,14,15	0.52	0	17,19,21	0.81	0
5	NAG	O	2002	1	14,14,15	0.40	0	17,19,21	1.48	3 (17%)
5	NAG	G	2001	1	14,14,15	0.46	0	17,19,21	1.40	1 (5%)
5	NAG	C	2003	1	14,14,15	0.58	0	17,19,21	1.63	3 (17%)
5	NAG	G	2002	1	14,14,15	0.47	0	17,19,21	1.73	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	E	2001	1	14,14,15	0.48	0	17,19,21	1.10	2 (11%)
5	NAG	A	2003	1	14,14,15	0.60	0	17,19,21	1.89	6 (35%)
5	NAG	K	2003	1	14,14,15	0.61	0	17,19,21	1.64	3 (17%)
5	NAG	I	2003	1	14,14,15	0.46	0	17,19,21	1.19	1 (5%)
5	NAG	K	2004	1	14,14,15	0.43	0	17,19,21	1.24	2 (11%)
5	NAG	Q	2004	1	14,14,15	0.53	0	17,19,21	1.31	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	2004	1	-	2/6/23/26	0/1/1/1
5	NAG	M	2003	1	-	2/6/23/26	0/1/1/1
5	NAG	O	2001	1	-	4/6/23/26	0/1/1/1
5	NAG	M	2004	1	-	0/6/23/26	0/1/1/1
5	NAG	O	2002	1	-	0/6/23/26	0/1/1/1
5	NAG	G	2001	1	-	2/6/23/26	0/1/1/1
5	NAG	C	2003	1	-	0/6/23/26	0/1/1/1
5	NAG	G	2002	1	-	2/6/23/26	0/1/1/1
5	NAG	E	2001	1	-	2/6/23/26	0/1/1/1
5	NAG	A	2003	1	-	3/6/23/26	0/1/1/1
5	NAG	K	2003	1	-	4/6/23/26	0/1/1/1
5	NAG	I	2003	1	-	2/6/23/26	0/1/1/1
5	NAG	K	2004	1	-	4/6/23/26	0/1/1/1
5	NAG	Q	2004	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	2002	NAG	C1-O5-C5	6.24	120.64	112.19
5	O	2001	NAG	C1-O5-C5	4.77	118.65	112.19
5	G	2001	NAG	C1-O5-C5	4.44	118.20	112.19
5	A	2003	NAG	C4-C3-C2	4.24	117.23	111.02
5	K	2003	NAG	O4-C4-C5	-4.06	99.22	109.30

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	2003	NAG	C8-C7-N2-C2
5	A	2003	NAG	O7-C7-N2-C2
5	E	2001	NAG	O7-C7-N2-C2
5	E	2004	NAG	C8-C7-N2-C2
5	E	2004	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	K	2004	NAG	3	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/334 (96%)	0.15	7 (2%) 62 56	62, 86, 117, 157	0
1	C	323/334 (96%)	0.41	23 (7%) 16 11	69, 102, 130, 161	0
1	E	323/334 (96%)	0.26	17 (5%) 26 20	67, 94, 121, 167	0
1	G	323/334 (96%)	0.21	3 (0%) 84 82	63, 85, 112, 148	0
1	I	323/334 (96%)	0.07	3 (0%) 84 82	46, 80, 100, 138	0
1	K	323/334 (96%)	0.30	14 (4%) 35 28	52, 93, 121, 164	0
1	M	323/334 (96%)	0.50	27 (8%) 11 7	79, 112, 143, 176	0
1	O	323/334 (96%)	0.29	18 (5%) 24 19	65, 102, 130, 154	0
1	Q	323/334 (96%)	0.07	5 (1%) 73 70	61, 82, 107, 135	0
2	B	173/182 (95%)	0.26	3 (1%) 70 66	54, 83, 115, 159	0
2	D	173/182 (95%)	0.33	4 (2%) 60 54	62, 91, 124, 171	0
2	F	173/182 (95%)	0.92	18 (10%) 6 4	57, 104, 165, 216	0
2	H	173/182 (95%)	0.75	15 (8%) 10 7	62, 101, 155, 194	0
2	J	173/182 (95%)	0.19	2 (1%) 79 76	58, 84, 120, 167	0
2	L	173/182 (95%)	0.62	11 (6%) 19 14	64, 101, 153, 167	0
2	N	173/182 (95%)	0.46	5 (2%) 51 45	69, 97, 135, 155	0
2	P	173/182 (95%)	0.56	6 (3%) 44 36	63, 91, 126, 181	0
2	R	173/182 (95%)	0.20	1 (0%) 89 88	61, 79, 106, 147	0
All	All	4464/4644 (96%)	0.33	182 (4%) 37 30	46, 91, 134, 216	0

The worst 5 of 182 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	9	PRO	10.1
2	P	173	ILE	8.6
2	F	140	PHE	8.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	9	PRO	7.0
2	H	173	ILE	6.8

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	U	2	14/15	0.80	0.35	135,145,154,158	0
3	NAG	S	2	14/15	0.81	0.24	121,139,156,167	0
3	NAG	X	2	14/15	0.82	0.19	144,161,171,177	0
4	MAN	Y	3	11/12	0.84	0.32	143,164,169,173	0
3	NAG	U	1	14/15	0.86	0.16	92,126,137,139	0
3	NAG	T	2	14/15	0.86	0.24	109,132,145,152	0
4	NAG	Y	2	14/15	0.87	0.20	122,141,155,162	0
3	NAG	Z	1	14/15	0.88	0.31	96,115,120,135	0
3	NAG	V	2	14/15	0.89	0.20	111,122,137,152	0
3	NAG	W	2	14/15	0.90	0.17	123,135,150,165	0
3	NAG	Z	2	14/15	0.91	0.22	119,142,154,163	0
3	NAG	T	1	14/15	0.92	0.12	82,103,115,116	0
3	NAG	X	1	14/15	0.93	0.10	91,109,123,143	0
3	NAG	V	1	14/15	0.93	0.12	68,84,96,98	0
4	NAG	Y	1	14/15	0.95	0.12	86,104,121,126	0
3	NAG	S	1	14/15	0.96	0.12	85,105,126,131	0
3	NAG	W	1	14/15	0.96	0.08	74,93,106,109	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.















Electron density around Chain W:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain X:**

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain Z:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain Y:**

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	K	2004	14/15	0.74	0.23	133,149,159,163	0
5	NAG	E	2004	14/15	0.81	0.22	119,136,150,151	0
5	NAG	K	2003	14/15	0.82	0.19	103,125,134,148	0
5	NAG	E	2001	14/15	0.84	0.30	103,118,138,158	0
5	NAG	M	2004	14/15	0.84	0.33	112,151,162,166	0
5	NAG	Q	2004	14/15	0.84	0.20	115,127,133,135	0
5	NAG	O	2001	14/15	0.86	0.18	75,105,120,122	0
5	NAG	A	2003	14/15	0.86	0.20	102,127,143,157	0
5	NAG	C	2003	14/15	0.88	0.31	115,124,136,149	0
5	NAG	M	2003	14/15	0.90	0.16	112,129,135,138	0
5	NAG	G	2001	14/15	0.91	0.15	86,106,114,116	0
5	NAG	G	2002	14/15	0.91	0.14	107,124,130,133	0
5	NAG	O	2002	14/15	0.91	0.12	92,112,118,125	0
5	NAG	I	2003	14/15	0.91	0.18	102,115,121,121	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.