



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 01:13 PM BST

PDB ID : 5QZK
Title : PanDDA analysis group deposition – Auto-refined data of Aar2/RNaseH for ground state model 35
Authors : Weiss, M.S.; Wollenhaupt, J.; Metz, A.; Barthel, T.; Lima, G.M.A.; Heine, A.; Mueller, U.; Klebe, G.
Deposited on : 2020-02-12
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	
2	B	308	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	2002	1283	335	372	12	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	300	2580	1654	421	485	20	0	9	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	-	linker	UNP P32357
B	167	SER	-	linker	UNP P32357
B	168	SER	-	linker	UNP P32357
B	169	SER	-	linker	UNP P32357
B	170	SER	-	linker	UNP P32357


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	17	Total 17	O 17	0	0
3	B	13	Total 13	O 13	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor 8



- Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.81Å 81.69Å 93.14Å 90.00° 108.28° 90.00°	Depositor
Resolution (Å)	44.65 – 1.73 44.61 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.0 (44.65-1.73) 99.1 (44.61-1.73)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.92 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.8.0238, PHENIX 1.16.3549	Depositor
R, R_{free}	0.267 , 0.269 0.272 , 0.285	Depositor DCC
R_{free} test set	3127 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	40.0	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4612	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/2049	0.76	0/2775
2	B	0.69	0/2651	0.77	0/3581
All	All	0.70	0/4700	0.77	0/6356

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2002	0	2029	8	0
2	B	2580	0	2448	20	0
3	A	17	0	0	0	0
3	B	13	0	0	0	0
All	All	4612	0	4477	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2062:GLU:O	1:A:2066:LYS:HG2	1.89	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1:MET:HB3	2:B:35:ASP:HA	1.73	0.69
2:B:77:LEU:HD21	2:B:79:LYS:HE3	1.80	0.64
2:B:287:ARG:O	2:B:291:ILE:HD13	2.02	0.59
2:B:2:ASN:HB3	2:B:62:CYS:SG	2.44	0.56
2:B:70:GLN:HB3	2:B:81:MET:HE1	1.88	0.56
2:B:16:GLY:HA3	2:B:45:HIS:CE1	2.41	0.56
2:B:2:ASN:CB	2:B:62:CYS:SG	2.94	0.56
2:B:188:ALA:HA	2:B:204:TYR:CD1	2.45	0.52
2:B:80:MET:HG2	2:B:137:PHE:CD2	2.46	0.51
2:B:280:SER:HB3	2:B:313:TYR:CE1	2.46	0.51
2:B:2:ASN:CG	2:B:62:CYS:HB3	2.32	0.50
1:A:1895:HIS:O	1:A:1898[A]:VAL:HG22	2.11	0.50
2:B:17:ILE:HD13	2:B:44[B]:ILE:HG13	1.94	0.50
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	1.94	0.49
2:B:17:ILE:HD13	2:B:44[B]:ILE:CG1	2.43	0.48
1:A:1962:ARG:O	1:A:2013:ARG:NH1	2.48	0.47
2:B:3:THR:HG21	2:B:93:VAL:HG13	1.96	0.46
1:A:1853:ASP:OD2	1:A:1855[B]:THR:HG23	2.15	0.46
1:A:2023:LYS:O	1:A:2027:LEU:HG	2.16	0.45
2:B:291:ILE:HG22	2:B:292:CYS:N	2.32	0.43
2:B:190:ARG:HD3	2:B:203[A]:TYR:CE2	2.54	0.42
2:B:258:LYS:HD2	2:B:258:LYS:H	1.85	0.41
1:A:1960[B]:GLU:HA	1:A:1960[B]:GLU:OE1	2.21	0.40
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.41	0.40
2:B:92:ILE:HD13	2:B:92:ILE:HA	1.94	0.40
2:B:77:LEU:N	2:B:77:LEU:HD23	2.37	0.40
2:B:80:MET:HG2	2:B:137:PHE:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/258 (96%)	242 (98%)	6 (2%)	0	100	100
2	B	306/308 (99%)	289 (94%)	17 (6%)	0	100	100
All	All	554/566 (98%)	531 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/233 (97%)	221 (98%)	5 (2%)	52	29
2	B	287/284 (101%)	287 (100%)	0	100	100
All	All	513/517 (99%)	508 (99%)	5 (1%)	86	63

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1962	ARG
1	A	1979[A]	MET
1	A	1979[B]	MET
1	A	1979[C]	MET
1	A	2027	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1869	ASN
1	A	1907	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	0.99	28 (11%) 4 5	26, 37, 75, 86	0
2	B	300/308 (97%)	1.03	40 (13%) 3 4	25, 39, 76, 125	0
All	All	537/566 (94%)	1.01	68 (12%) 3 4	25, 38, 76, 125	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	MET	9.1
2	B	52	SER	8.2
1	A	2028	SER	8.1
1	A	1833	GLY	7.8
2	B	53	SER	7.7
1	A	2063	TYR	7.7
2	B	109	ASP	6.7
2	B	170	SER	6.6
1	A	2027	LEU	6.3
2	B	173	ALA	5.8
2	B	108	ILE	5.7
2	B	279	TYR	5.7
2	B	22	PHE	5.6
2	B	54[A]	MET	5.5
1	A	2064	GLY	5.4
1	A	2046	GLU	5.4
1	A	2060	LEU	5.2
2	B	172	PRO	4.9
1	A	1962	ARG	4.8
2	B	171	ASP	4.6
1	A	2066	LYS	4.5
2	B	275	LEU	4.3
1	A	1836	ASN	3.8
1	A	2068	ASN	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	133	ASP	3.6
2	B	174	HIS	3.5
1	A	1835	MET	3.3
2	B	84	ARG	3.2
1	A	1953	ASN	3.2
2	B	41[A]	VAL	3.2
1	A	1840	TYR	3.1
2	B	203[A]	TYR	3.1
1	A	1866	PHE	3.0
2	B	280	SER	2.9
1	A	1837	SER	2.8
2	B	247	ILE	2.8
1	A	1839	ASN	2.8
2	B	129	ILE	2.7
2	B	110	GLU	2.7
2	B	136	GLN	2.7
2	B	101	MET	2.6
2	B	316	LEU	2.6
2	B	106	PRO	2.6
2	B	9	ALA	2.6
2	B	151	GLU	2.5
1	A	2067	TYR	2.5
1	A	2049	ILE	2.5
1	A	1961	LEU	2.5
2	B	195	MET	2.4
2	B	44[A]	ILE	2.4
2	B	293	LEU	2.4
2	B	313	TYR	2.4
2	B	224	LEU	2.3
2	B	278	GLN	2.2
2	B	277	GLU	2.2
1	A	1996	LEU	2.2
1	A	2051	ILE	2.2
2	B	134	GLU	2.2
1	A	1870	VAL	2.2
2	B	308	ILE	2.2
2	B	114	TRP	2.2
1	A	1920	LEU	2.1
2	B	211[A]	GLN	2.1
2	B	121	VAL	2.1
1	A	2061	THR	2.1
1	A	2057[A]	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2065	ARG	2.0
1	A	1989	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.