



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 04:50 am BST

PDB ID : 4TV7
Title : Crystal structure of Bacillus subtilis GabR at 2.05 Angstroms resolution
Authors : Goto, M.; Okuda, K.; Yoshimura, T.
Deposited on : 2014-06-26
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	
1	B	479	
1	C	479	
1	D	479	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 14499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HTH-type transcriptional regulatory protein GabR.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	432	Total 3344	C 2135	N 570	O 621	P 1	S 17	0	0	0
1	B	430	Total 3373	C 2152	N 575	O 629	P 1	S 16	0	0	0
1	C	452	Total 3623	C 2303	N 623	O 679	P 1	S 17	0	0	0
1	D	449	Total 3596	C 2286	N 610	O 683	P 1	S 16	0	0	0


- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	97	Total 97	O 97	0	0
2	B	120	Total 120	O 120	0	0
2	C	181	Total 181	O 181	0	0
2	D	165	Total 165	O 165	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HTH-type transcriptional regulatory protein GabR






• Molecule 1: HTH-type transcriptional regulatory protein GabR



• Molecule 1: HTH-type transcriptional regulatory protein GabR



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.83Å 101.56Å 212.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.67 – 2.05 47.67 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.3 (47.67-2.05) 97.3 (47.67-2.05)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.53 (at 2.05Å)	Xtrriage
Refinement program	REFMAC refmac_5.7.0032, REFMAC, REFMAC5	Depositor
R, R_{free}	0.231 , 0.284 0.230 , 0.283	Depositor DCC
R_{free} test set	6564 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	43.3	Xtrriage
Anisotropy	0.473	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.013 for k,h,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14499	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3380	0.66	3/4573 (0.1%)
1	B	0.46	0/3411	0.66	1/4606 (0.0%)
1	C	0.55	0/3666	0.75	3/4947 (0.1%)
1	D	0.54	0/3638	0.74	3/4910 (0.1%)
All	All	0.50	0/14095	0.71	10/19036 (0.1%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	155	ARG	NE-CZ-NH2	-11.63	114.48	120.30
1	C	155	ARG	NE-CZ-NH1	10.31	125.46	120.30
1	D	211	LEU	CA-CB-CG	6.83	131.02	115.30
1	A	211	LEU	CA-CB-CG	6.45	130.12	115.30
1	C	211	LEU	CA-CB-CG	6.43	130.09	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3344	0	3211	63	0
1	B	3373	0	3265	58	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3623	0	3576	55	0
1	D	3596	0	3536	46	0
2	A	97	0	0	2	0
2	B	120	0	0	6	0
2	C	181	0	0	14	0
2	D	165	0	0	6	0
All	All	14499	0	13588	212	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 212 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:312:LLP:H6	2:C:666:HOH:O	1.48	1.11
1:A:16:ILE:HG21	1:A:51:VAL:HG11	1.42	0.99
1:D:328:GLU:HG3	2:D:646:HOH:O	1.62	0.97
1:A:51:VAL:HG12	2:A:576:HOH:O	1.70	0.90
1:D:80:GLU:O	1:D:81:GLU:HB3	1.73	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	417/479 (87%)	377 (90%)	33 (8%)	7 (2%)	9 2
1	B	413/479 (86%)	384 (93%)	21 (5%)	8 (2%)	8 2
1	C	441/479 (92%)	418 (95%)	14 (3%)	9 (2%)	7 1
1	D	438/479 (91%)	411 (94%)	20 (5%)	7 (2%)	9 2
All	All	1709/1916 (89%)	1590 (93%)	88 (5%)	31 (2%)	8 2

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	121	HIS
1	A	390	VAL
1	B	407	THR
1	C	11	GLU
1	C	139	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	335/424 (79%)	310 (92%)	25 (8%)	13 6
1	B	347/424 (82%)	324 (93%)	23 (7%)	16 9
1	C	386/424 (91%)	366 (95%)	20 (5%)	23 14
1	D	383/424 (90%)	351 (92%)	32 (8%)	11 4
All	All	1451/1696 (86%)	1351 (93%)	100 (7%)	15 8

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	403	THR
1	C	145	MET
1	D	330	LEU
1	B	415	LEU
1	C	41	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	370	HIS
1	C	147	HIS
1	D	422	GLN
1	B	417	HIS
1	C	121	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	D	312	1	23,24,25	3.12	6 (26%)	25,32,34	1.69	7 (28%)
1	LLP	C	312	1	23,24,25	2.95	5 (21%)	25,32,34	1.48	5 (20%)
1	LLP	B	312	1	23,24,25	2.86	5 (21%)	25,32,34	1.67	6 (24%)
1	LLP	A	312	1	23,24,25	2.91	5 (21%)	25,32,34	1.43	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	D	312	1	-	7/16/17/19	0/1/1/1
1	LLP	C	312	1	-	7/16/17/19	0/1/1/1
1	LLP	B	312	1	-	8/16/17/19	0/1/1/1
1	LLP	A	312	1	-	10/16/17/19	0/1/1/1

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	312	LLP	C3-C2	8.16	1.49	1.40
1	D	312	LLP	C3-C2	7.94	1.48	1.40
1	D	312	LLP	C4-C5	7.73	1.51	1.42
1	C	312	LLP	C4-C5	7.66	1.51	1.42
1	A	312	LLP	C4-C5	7.37	1.51	1.42

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	312	LLP	OP4-C5'-C5	4.65	118.20	109.35
1	D	312	LLP	OP4-C5'-C5	4.20	117.36	109.35
1	B	312	LLP	C4-C4'-NZ	-3.52	108.13	124.31
1	D	312	LLP	C4-C4'-NZ	-3.51	108.21	124.31
1	C	312	LLP	C4-C4'-NZ	-3.21	109.58	124.31

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	312	LLP	C5'-OP4-P-OP2
1	D	312	LLP	C5'-OP4-P-OP3
1	C	312	LLP	C5'-OP4-P-OP2
1	C	312	LLP	C5'-OP4-P-OP3
1	B	312	LLP	C4-C4'-NZ-CE

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	312	LLP	1	0
1	C	312	LLP	4	0
1	B	312	LLP	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	431/479 (89%)	0.78	56 (12%) 3 2	32, 58, 98, 114	0
1	B	429/479 (89%)	0.73	46 (10%) 6 6	35, 56, 84, 115	1 (0%)
1	C	451/479 (94%)	0.48	22 (4%) 29 31	30, 45, 73, 99	1 (0%)
1	D	448/479 (93%)	0.40	19 (4%) 36 39	29, 46, 76, 105	1 (0%)
All	All	1759/1916 (91%)	0.59	143 (8%) 12 12	29, 50, 86, 115	3 (0%)

The worst 5 of 143 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	390	VAL	7.6
1	A	419	ALA	7.1
1	C	142	LEU	6.9
1	D	316	PRO	6.7
1	A	386	PHE	6.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	D	312	24/25	0.78	0.29	51,75,83,85	0
1	LLP	A	312	24/25	0.78	0.26	64,85,95,103	0
1	LLP	B	312	24/25	0.80	0.22	50,73,83,87	0
1	LLP	C	312	24/25	0.80	0.26	44,72,82,87	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.