



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 4, 2024 – 12:49 pm GMT

PDB ID : 4UHV
Title : The structure of VgrG1, the needle tip of the bacterial Type VI Secretion System
Authors : Spinola-Amilibia, M.; Davo-Siguero, I.; Ruiz, F.M.; Santillana, E.; Medrano, F.J.; Romero, A.
Deposited on : 2015-03-25
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	651	 3% 55% 35% 7% ..
1	B	651	 % 60% 33% 5% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	B	1657	-	-	X	-

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 10911 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VGRG1, VALINE-GLYCINE REPEAT PROTEIN G1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	644	5107	3194	933	964	7	9	0	3	0
1	B	645	5109	3194	930	969	7	9	0	3	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	644	LEU	-	expression tag	UNP Q9I741
A	645	GLU	-	expression tag	UNP Q9I741
A	646	HIS	-	expression tag	UNP Q9I741
A	647	HIS	-	expression tag	UNP Q9I741
A	648	HIS	-	expression tag	UNP Q9I741
A	649	HIS	-	expression tag	UNP Q9I741
A	650	HIS	-	expression tag	UNP Q9I741
A	651	HIS	-	expression tag	UNP Q9I741
B	644	LEU	-	expression tag	UNP Q9I741
B	645	GLU	-	expression tag	UNP Q9I741
B	646	HIS	-	expression tag	UNP Q9I741
B	647	HIS	-	expression tag	UNP Q9I741
B	648	HIS	-	expression tag	UNP Q9I741
B	649	HIS	-	expression tag	UNP Q9I741
B	650	HIS	-	expression tag	UNP Q9I741
B	651	HIS	-	expression tag	UNP Q9I741

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	13	Total	Cl	0	0
			13	13		
2	B	25	Total	Cl	0	0
			25	25		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	6	Total 6	Na 6	0	0
3	B	7	Total 7	Na 7	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	268	Total 268	O 268	0	0
4	B	376	Total 376	O 376	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: VGRG1, VALINE-GLYCINE REPEAT PROTEIN G1



- Molecule 1: VGRG1, VALINE-GLYCINE REPEAT PROTEIN G1





4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	78.62Å 78.62Å 430.32Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	71.73 – 2.00 71.72 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.0 (71.73-2.00) 94.0 (71.72-2.00)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.223 , 0.248 0.225 , 0.249	Depositor DCC
R_{free} test set	4702 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.284	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 66.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.34$, $\langle L^2 \rangle = 0.17$	Xtriage
Estimated twinning fraction	0.480 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.500 for H, K, L 0.500 for K, H, -L	Depositor
Outliers	0 of 94822 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	10911	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/5228	0.80	4/7069 (0.1%)
1	B	0.57	0/5230	0.85	5/7074 (0.1%)
All	All	0.54	0/10458	0.82	9/14143 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	9
1	B	0	5
All	All	0	14

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	460	LEU	C-N-CD	-15.80	85.84	120.60
1	B	258	TYR	C-N-CD	-6.73	105.79	120.60
1	A	18	LEU	CA-CB-CG	6.24	129.66	115.30
1	B	439	ASP	CB-CG-OD2	-5.71	113.16	118.30
1	A	486	ARG	NE-CZ-NH1	5.58	123.09	120.30

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	110	GLN	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	246	ARG	Peptide
1	A	258	TYR	Peptide
1	A	267	ASP	Peptide
1	A	3	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5107	0	4940	218	3
1	B	5109	0	4932	189	2
2	A	13	0	0	3	0
2	B	25	0	0	7	0
3	A	6	0	0	0	0
3	B	7	0	0	0	0
4	A	268	0	0	30	1
4	B	376	0	0	28	0
All	All	10911	0	9872	408	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 408 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:397:HIS:HD2	4:B:2179:HOH:O	1.15	1.24
1:B:258:TYR:CD2	1:B:259:PRO:HD2	1.88	1.09
1:A:11:CYS:SG	1:A:12:PRO:HD3	1.93	1.08
1:A:572:CYS:HA	4:A:2224:HOH:O	1.53	1.05
1:A:522:LYS:HE2	1:A:524:ILE:HD11	1.40	1.02

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:564:ALA:O	4:A:2224:HOH:O[3_765]	1.76	0.44
1:A:503:ASN:ND2	1:A:511:ASP:O[3_765]	1.99	0.21
1:A:45:ASN:OD1	1:A:66:SER:OG[2_755]	2.04	0.16
1:B:465:THR:OG1	1:B:474:SER:OG[2_755]	2.12	0.08
1:B:523:SER:OG	1:B:531:ARG:NH2[2_755]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	645/651 (99%)	519 (80%)	103 (16%)	23 (4%)	3	1
1	B	646/651 (99%)	538 (83%)	88 (14%)	20 (3%)	4	1
All	All	1291/1302 (99%)	1057 (82%)	191 (15%)	43 (3%)	4	1

5 of 43 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	LEU
1	A	63	PRO
1	A	64	GLY
1	A	234	PRO
1	A	249	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	539/534 (101%)	445 (83%)	94 (17%)	2	1
1	B	540/534 (101%)	477 (88%)	63 (12%)	5	3
All	All	1079/1068 (101%)	922 (85%)	157 (15%)	3	1

5 of 157 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	205	ARG
1	B	520	ARG
1	B	273	ARG
1	B	391	HIS
1	B	570	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	71	HIS
1	B	266	GLN
1	B	586	ASN
1	B	243	ASN
1	B	339	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 51 ligands modelled in this entry, 51 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	635/651 (97%)	0.06	21 (3%) 46 45	15, 41, 63, 78	0
1	B	636/651 (97%)	-0.22	5 (0%) 86 85	10, 33, 54, 63	0
All	All	1271/1302 (97%)	-0.08	26 (2%) 65 63	10, 37, 59, 78	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	336	ALA	4.3
1	A	333	SER	3.6
1	A	196	PRO	3.6
1	B	641	ALA	3.3
1	B	9	VAL	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	B	1656	1/1	0.85	0.07	59,59,59,59	0
2	CL	B	1707	1/1	0.89	0.10	75,75,75,75	0
2	CL	B	1654	1/1	0.94	0.05	52,52,52,52	0
2	CL	B	1647	1/1	0.94	0.11	63,63,63,63	0
2	CL	B	1660	1/1	0.94	0.08	50,50,50,50	0
2	CL	B	1648	1/1	0.94	0.18	70,70,70,70	0
3	NA	A	1711	1/1	0.94	0.11	33,33,33,33	0
3	NA	B	1663	1/1	0.94	0.12	60,60,60,60	0
3	NA	B	1711	1/1	0.94	0.15	48,48,48,48	0
3	NA	B	1712	1/1	0.94	0.14	40,40,40,40	0
2	CL	B	1655	1/1	0.95	0.05	68,68,68,68	0
2	CL	A	1704	1/1	0.95	0.05	42,42,42,42	0
2	CL	A	1649	1/1	0.96	0.06	43,43,43,43	0
3	NA	A	1655	1/1	0.96	0.07	34,34,34,34	0
2	CL	A	1650	1/1	0.96	0.05	57,57,57,57	0
2	CL	A	1647	1/1	0.96	0.10	44,44,44,44	0
2	CL	B	1704	1/1	0.96	0.04	48,48,48,48	0
2	CL	B	1705	1/1	0.96	0.07	42,42,42,42	0
3	NA	B	1713	1/1	0.96	0.10	43,43,43,43	0
2	CL	B	1661	1/1	0.97	0.05	39,39,39,39	0
2	CL	B	1702	1/1	0.97	0.07	35,35,35,35	0
3	NA	B	1710	1/1	0.97	0.07	32,32,32,32	0
3	NA	A	1653	1/1	0.97	0.09	35,35,35,35	0
3	NA	A	1654	1/1	0.97	0.18	41,41,41,41	0
2	CL	B	1649	1/1	0.97	0.05	55,55,55,55	0
2	CL	B	1706	1/1	0.98	0.07	46,46,46,46	0
2	CL	A	1701	1/1	0.98	0.04	40,40,40,40	0
2	CL	B	1708	1/1	0.98	0.03	50,50,50,50	0
3	NA	A	1652	1/1	0.98	0.06	34,34,34,34	0
2	CL	A	1703	1/1	0.98	0.04	51,51,51,51	0
2	CL	B	1658	1/1	0.98	0.08	46,46,46,46	0
2	CL	B	1659	1/1	0.98	0.08	53,53,53,53	0
2	CL	A	1646	1/1	0.98	0.11	41,41,41,41	0
2	CL	B	1651	1/1	0.98	0.09	44,44,44,44	0
2	CL	B	1652	1/1	0.98	0.05	43,43,43,43	0
2	CL	B	1703	1/1	0.98	0.07	41,41,41,41	0
2	CL	B	1653	1/1	0.98	0.04	43,43,43,43	0
2	CL	B	1646	1/1	0.98	0.08	49,49,49,49	0
2	CL	A	1700	1/1	0.99	0.09	38,38,38,38	0
2	CL	B	1700	1/1	0.99	0.07	26,26,26,26	0
2	CL	B	1701	1/1	0.99	0.07	46,46,46,46	0
2	CL	A	1645	1/1	0.99	0.10	31,31,31,31	0
3	NA	A	1710	1/1	0.99	0.10	36,36,36,36	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	A	1702	1/1	0.99	0.16	38,38,38,38	0
3	NA	B	1662	1/1	0.99	0.07	20,20,20,20	0
2	CL	A	1648	1/1	0.99	0.10	37,37,37,37	0
2	CL	B	1657	1/1	0.99	0.08	42,42,42,42	0
2	CL	B	1650	1/1	0.99	0.04	37,37,37,37	0
2	CL	A	1651	1/1	0.99	0.11	45,45,45,45	0
2	CL	A	1705	1/1	0.99	0.04	40,40,40,40	0
3	NA	B	1714	1/1	0.99	0.16	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.