



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 25, 2020 – 04:24 PM BST

PDB ID : 3VA7  
Title : Crystal structure of the Kluyveromyces lactis Urea Carboxylase  
Authors : Fan, C.; Xiang, S.  
Deposited on : 2011-12-29  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1236	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	URE	A	1902	-	-	X	-
4	GOL	A	1905	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
4	GOL	A	1906	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9268 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called KLLA0E08119p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1130	8829	5608	1499	1686	36	0	0	0

There are 23 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	594	MET	-	EXPRESSION TAG	UNP Q6CP22
A	595	GLY	-	EXPRESSION TAG	UNP Q6CP22
A	596	SER	-	EXPRESSION TAG	UNP Q6CP22
A	597	SER	-	EXPRESSION TAG	UNP Q6CP22
A	598	HIS	-	EXPRESSION TAG	UNP Q6CP22
A	599	HIS	-	EXPRESSION TAG	UNP Q6CP22
A	600	HIS	-	EXPRESSION TAG	UNP Q6CP22
A	601	HIS	-	EXPRESSION TAG	UNP Q6CP22
A	602	HIS	-	EXPRESSION TAG	UNP Q6CP22
A	603	HIS	-	EXPRESSION TAG	UNP Q6CP22
A	604	SER	-	EXPRESSION TAG	UNP Q6CP22
A	605	SER	-	EXPRESSION TAG	UNP Q6CP22
A	606	GLY	-	EXPRESSION TAG	UNP Q6CP22
A	607	LEU	-	EXPRESSION TAG	UNP Q6CP22
A	608	VAL	-	EXPRESSION TAG	UNP Q6CP22
A	609	PRO	-	EXPRESSION TAG	UNP Q6CP22
A	610	ARG	-	EXPRESSION TAG	UNP Q6CP22
A	611	GLY	-	EXPRESSION TAG	UNP Q6CP22
A	612	SER	-	EXPRESSION TAG	UNP Q6CP22
A	613	HIS	-	EXPRESSION TAG	UNP Q6CP22
A	614	MET	-	EXPRESSION TAG	UNP Q6CP22
A	615	ALA	-	EXPRESSION TAG	UNP Q6CP22
A	616	SER	-	EXPRESSION TAG	UNP Q6CP22

- Molecule 2 is 5-(HEXAHYDRO-2-OXO-1H-THIENO[3,4-D]IMIDAZOL-6-YL)PENTANAL (three-letter code: BTI) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	15	10	2	2	1	0	0

- Molecule 3 is UREA (three-letter code: URE) (formula:  $\text{CH}_4\text{N}_2\text{O}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	4	1	2	1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $\text{C}_3\text{H}_8\text{O}_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	377	Total O 377 377	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	126.66Å 126.66Å 217.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.60 47.73 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.1 (50.00-2.60) 99.1 (47.73-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.75 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.187 , 0.255 0.184 , 0.249	Depositor DCC
$R_{free}$ test set	2790 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.0	Xtrriage
Anisotropy	0.512	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 42.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9268	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BTI, URE, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.94	4/9017 (0.0%)	0.94	13/12220 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1311	GLU	CG-CD	5.92	1.60	1.51
1	A	1114	VAL	CB-CG1	-5.78	1.40	1.52
1	A	1612	TYR	CG-CD2	5.55	1.46	1.39
1	A	1564	ASP	CB-CG	5.07	1.62	1.51

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1467	ARG	NE-CZ-NH2	-8.00	116.30	120.30
1	A	1467	ARG	NE-CZ-NH1	6.52	123.56	120.30
1	A	989	ASP	CB-CG-OD1	6.31	123.98	118.30
1	A	1438	ASP	N-CA-CB	-6.11	99.60	110.60
1	A	1203	MET	CG-SD-CE	-5.97	90.64	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8829	0	8753	213	0
2	A	15	0	15	0	0
3	A	4	0	4	3	0
4	A	42	0	56	22	0
5	A	1	0	0	0	0
6	A	377	0	0	17	0
All	All	9268	0	8828	215	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 215 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:914:ARG:HA	4:A:1905:GOL:H32	1.27	1.12
1:A:1025:GLN:HG3	6:A:2003:HOH:O	1.52	1.04
1:A:1565:ARG:HD3	6:A:2069:HOH:O	1.68	0.94
1:A:1459:ASN:HD21	1:A:1460:ARG:HH11	1.15	0.93
1:A:1553:ASN:HD22	1:A:1639:LEU:H	1.22	0.88

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1124/1236 (91%)	1046 (93%)	71 (6%)	7 (1%)	<b>25</b> 47

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1166	ARG
1	A	1438	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1711	GLU
1	A	891	CYS
1	A	1003	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	955/1040 (92%)	859 (90%)	96 (10%)	<b>7</b> <b>14</b>

5 of 96 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1074	LEU
1	A	1224	GLN
1	A	1736	GLN
1	A	1079	ASN
1	A	1150	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1318	ASN
1	A	1389	GLN
1	A	1638	GLN
1	A	1278	GLN
1	A	1315	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	1908	-	5,5,5	0.98	0	5,5,5	1.01	0
4	GOL	A	1907	-	5,5,5	0.48	0	5,5,5	0.47	0
4	GOL	A	1904	-	5,5,5	0.57	0	5,5,5	0.90	0
4	GOL	A	1903	-	5,5,5	0.53	0	5,5,5	0.93	0
4	GOL	A	1909	-	5,5,5	0.59	0	5,5,5	0.61	0
2	BTI	A	1901	1	16,16,16	1.72	3 (18%)	21,21,21	2.58	7 (33%)
4	GOL	A	1905	-	5,5,5	0.79	0	5,5,5	1.21	0
4	GOL	A	1906	-	5,5,5	0.35	0	5,5,5	1.00	0
3	URE	A	1902	-	3,3,3	0.60	0	3,3,3	2.38	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1908	-	-	2/4/4/4	-
4	GOL	A	1907	-	-	1/4/4/4	-
4	GOL	A	1904	-	-	1/4/4/4	-
4	GOL	A	1903	-	-	0/4/4/4	-
4	GOL	A	1909	-	-	0/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTI	A	1901	1	-	1/5/27/27	0/2/2/2
4	GOL	A	1905	-	-	4/4/4/4	-
4	GOL	A	1906	-	-	2/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1901	BTI	O3-C3	4.21	1.32	1.23
2	A	1901	BTI	C2-S1	-3.33	1.77	1.82
2	A	1901	BTI	C3-N3	-2.51	1.31	1.35

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1901	BTI	C6-C5-N3	-8.51	102.21	113.03
2	A	1901	BTI	C2-C4-N2	-4.98	108.66	113.13
2	A	1901	BTI	N2-C3-N3	3.69	112.22	108.76
3	A	1902	URE	N2-C-N1	-3.40	111.44	117.82
2	A	1901	BTI	C6-C5-C4	2.40	110.74	108.66

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1908	GOL	C1-C2-C3-O3
4	A	1908	GOL	O2-C2-C3-O3
4	A	1905	GOL	O1-C1-C2-C3
4	A	1905	GOL	C1-C2-C3-O3
4	A	1906	GOL	C1-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1908	GOL	2	0
4	A	1904	GOL	1	0
4	A	1909	GOL	1	0
4	A	1905	GOL	12	0
4	A	1906	GOL	6	0
3	A	1902	URE	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1130/1236 (91%)	-0.42	14 (1%) 79 76	18, 37, 65, 105	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1227	LEU	4.3
1	A	1829	VAL	3.4
1	A	1065	ASP	3.2
1	A	1755	GLU	3.1
1	A	754	PRO	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	1908	6/6	0.85	0.29	46,53,54,55	0
4	GOL	A	1904	6/6	0.85	0.24	66,70,72,72	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	1905	6/6	0.90	0.33	48,50,51,52	0
4	GOL	A	1909	6/6	0.91	0.17	54,55,57,58	0
4	GOL	A	1906	6/6	0.95	0.16	52,53,54,54	0
4	GOL	A	1903	6/6	0.96	0.17	43,48,50,54	0
3	URE	A	1902	4/4	0.97	0.22	39,39,40,40	0
5	NA	A	1910	1/1	0.98	0.14	62,62,62,62	0
4	GOL	A	1907	6/6	0.98	0.14	34,37,40,43	0
2	BTI	A	1901	15/15	0.99	0.20	18,24,26,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.