



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 11:10 pm BST

PDB ID : 4XA9
Title : Crystal structure of the complex between the N-terminal domain of RavJ and LegL1 from Legionella pneumophila str. Philadelphia
Authors : Stogios, P.J.; Cuff, M.E.; Nocek, B.; Evdokimova, E.; Di Leo, R.; Yim, V.; Savchenko, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2014-12-13
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	81% 17%
1	B	297	77% 6% 17%
1	C	297	80% 17%
1	D	297	81% 17%
1	E	297	% 79% 17%
1	F	297	79% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	297	 82% 16%
1	H	297	 81% 17%
2	a	231	 97% 4%
2	b	231	 97% 4%
2	c	231	 96% 4%
2	d	231	 97% 2%
2	e	231	 97% 2%
2	f	231	 97% 2%
2	g	231	 97% 1%
2	h	231	 98% 2%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 34121 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gala protein type 1, 3 or 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	247	Total 1916	C 1209	N 319	O 384	S 4	0	10	0
1	B	247	Total 1907	C 1200	N 317	O 386	S 4	0	8	0
1	C	247	Total 1889	C 1189	N 316	O 379	S 5	0	3	0
1	D	247	Total 1897	C 1195	N 316	O 382	S 4	0	6	0
1	E	247	Total 1898	C 1192	N 317	O 385	S 4	0	6	0
1	F	247	Total 1904	C 1198	N 319	O 383	S 4	0	6	0
1	G	248	Total 1900	C 1195	N 319	O 382	S 4	0	4	0
1	H	247	Total 1894	C 1188	N 318	O 384	S 4	0	5	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q5ZWY8
B	0	GLY	-	expression tag	UNP Q5ZWY8
C	0	GLY	-	expression tag	UNP Q5ZWY8
D	0	GLY	-	expression tag	UNP Q5ZWY8
E	0	GLY	-	expression tag	UNP Q5ZWY8
F	0	GLY	-	expression tag	UNP Q5ZWY8
G	0	GLY	-	expression tag	UNP Q5ZWY8
H	0	GLY	-	expression tag	UNP Q5ZWY8

- Molecule 2 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	a	228	Total	C	N	O	S	Se	0	3	0
			1796	1139	312	340	2	3			
2	b	227	Total	C	N	O	S	Se	0	1	0
			1779	1128	310	337	2	2			
2	c	224	Total	C	N	O	S	Se	0	3	0
			1779	1132	309	334	2	2			
2	d	228	Total	C	N	O	S	Se	0	5	0
			1808	1149	313	341	2	3			
2	e	227	Total	C	N	O	S	Se	0	2	0
			1787	1134	311	337	2	3			
2	f	226	Total	C	N	O	S	Se	0	7	0
			1810	1151	314	340	2	3			
2	g	225	Total	C	N	O	S	Se	0	4	0
			1782	1132	310	335	2	3			
2	h	229	Total	C	N	O	S	Se	0	2	0
			1794	1137	312	340	2	3			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	0	GLY	-	expression tag	UNP Q5ZWY9
b	0	GLY	-	expression tag	UNP Q5ZWY9
c	0	GLY	-	expression tag	UNP Q5ZWY9
d	0	GLY	-	expression tag	UNP Q5ZWY9
e	0	GLY	-	expression tag	UNP Q5ZWY9
f	0	GLY	-	expression tag	UNP Q5ZWY9
g	0	GLY	-	expression tag	UNP Q5ZWY9
h	0	GLY	-	expression tag	UNP Q5ZWY9

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	284	Total	O	0	3
			287	287		
3	a	230	Total	O	0	4
			234	234		
3	B	288	Total	O	0	7
			295	295		
3	b	243	Total	O	0	5
			248	248		
3	C	326	Total	O	0	6
			332	332		
3	c	256	Total	O	0	5
			261	261		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	300	Total 306	O 306	0	6
3	d	249	Total 254	O 254	0	5
3	E	288	Total 294	O 294	0	6
3	e	306	Total 311	O 311	0	5
3	F	297	Total 301	O 301	0	4
3	f	338	Total 342	O 342	0	4
3	G	293	Total 295	O 295	0	2
3	g	273	Total 274	O 274	0	1
3	H	260	Total 262	O 262	0	2
3	h	281	Total 285	O 285	0	4

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gala protein type 1, 3 or 4

Chain A: 



- Molecule 1: Gala protein type 1, 3 or 4

Chain B: 



- Molecule 1: Gala protein type 1, 3 or 4

Chain C: 




- Molecule 1: Gala protein type 1, 3 or 4


Chain D: 




- Molecule 1: Gala protein type 1, 3 or 4




- Molecule 1: Gala protein type 1, 3 or 4




- Molecule 1: Gala protein type 1, 3 or 4



- Molecule 1: Gala protein type 1, 3 or 4



- Molecule 2: Uncharacterized protein



- Molecule 2: Uncharacterized protein






- Molecule 2: Uncharacterized protein




- Molecule 2: Uncharacterized protein



- Molecule 2: Uncharacterized protein




- Molecule 2: Uncharacterized protein



- Molecule 2: Uncharacterized protein



- Molecule 2: Uncharacterized protein



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	219.61Å 120.89Å 149.29Å 90.00° 92.05° 90.00°	Depositor
Resolution (Å)	29.94 – 2.00 29.94 – 1.99	Depositor EDS
% Data completeness (in resolution range)	86.6 (29.94-2.00) 95.6 (29.94-1.99)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.22 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.158 , 0.195 0.158 , 0.194	Depositor DCC
R_{free} test set	12932 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtrriage
Anisotropy	0.067	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 62.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.001 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.004 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.010 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.003 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.027 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	34121	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.81 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6723e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1964	0.47	1/2663 (0.0%)
1	B	0.33	0/1952	0.45	0/2650
1	C	0.30	0/1919	0.45	0/2603
1	D	0.28	0/1936	0.44	0/2627
1	E	0.28	0/1934	0.43	0/2625
1	F	0.29	0/1941	0.44	0/2635
1	G	0.30	0/1934	0.45	0/2626
1	H	0.28	0/1930	0.44	0/2620
2	a	0.37	0/1850	0.49	0/2516
2	b	0.35	0/1827	0.46	0/2486
2	c	0.32	0/1833	0.46	0/2492
2	d	0.36	0/1868	0.47	0/2541
2	e	0.40	0/1838	0.50	0/2500
2	f	0.36	0/1876	0.47	0/2551
2	g	0.37	0/1838	0.47	0/2497
2	h	0.40	0/1844	0.48	0/2506
All	All	0.34	0/30284	0.46	1/41138 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	85	TRP	CA-CB-CG	-5.49	103.27	113.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1916	0	1982	4	0
1	B	1907	0	1951	10	0
1	C	1889	0	1931	3	0
1	D	1897	0	1946	3	0
1	E	1898	0	1934	6	0
1	F	1904	0	1942	9	0
1	G	1900	0	1936	3	0
1	H	1894	0	1926	4	0
2	a	1796	0	1778	0	0
2	b	1779	0	1753	0	0
2	c	1779	0	1761	0	0
2	d	1808	0	1799	0	0
2	e	1787	0	1768	0	0
2	f	1810	0	1803	0	0
2	g	1782	0	1772	0	0
2	h	1794	0	1773	0	0
3	A	287	0	0	0	0
3	B	295	0	0	2	0
3	C	332	0	0	0	0
3	D	306	0	0	1	0
3	E	294	0	0	0	0
3	F	301	0	0	2	0
3	G	295	0	0	1	0
3	H	262	0	0	1	0
3	a	234	0	0	0	0
3	b	248	0	0	0	0
3	c	261	0	0	0	0
3	d	254	0	0	0	0
3	e	311	0	0	0	0
3	f	342	0	0	0	0
3	g	274	0	0	0	0
3	h	285	0	0	0	0
All	All	34121	0	29755	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:107:VAL:HG11	1:F:126:ILE:HD13	1.71	0.70
1:A:167:ILE:HD12	1:A:191:ILE:HD12	1.86	0.57
1:B:5:LEU:HD21	1:E:5:LEU:HD21	1.89	0.55
1:A:5:LEU:HD21	1:F:5:LEU:HD21	1.91	0.52
1:H:60:LYS:HD3	3:H:560:HOH:O	2.10	0.52

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/297 (86%)	244 (96%)	11 (4%)	0	100	100
1	B	253/297 (85%)	243 (96%)	10 (4%)	0	100	100
1	C	248/297 (84%)	239 (96%)	8 (3%)	1 (0%)	34	30
1	D	251/297 (84%)	240 (96%)	11 (4%)	0	100	100
1	E	251/297 (84%)	243 (97%)	7 (3%)	1 (0%)	34	30
1	F	251/297 (84%)	241 (96%)	10 (4%)	0	100	100
1	G	250/297 (84%)	239 (96%)	11 (4%)	0	100	100
1	H	250/297 (84%)	240 (96%)	10 (4%)	0	100	100
2	a	229/231 (99%)	226 (99%)	3 (1%)	0	100	100
2	b	226/231 (98%)	224 (99%)	2 (1%)	0	100	100
2	c	225/231 (97%)	224 (100%)	1 (0%)	0	100	100
2	d	231/231 (100%)	227 (98%)	4 (2%)	0	100	100
2	e	227/231 (98%)	224 (99%)	3 (1%)	0	100	100
2	f	231/231 (100%)	229 (99%)	2 (1%)	0	100	100
2	g	227/231 (98%)	222 (98%)	5 (2%)	0	100	100
2	h	229/231 (99%)	227 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	3834/4224 (91%)	3732 (97%)	100 (3%)	2 (0%)	51	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	123	TYR
1	C	123	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/260 (85%)	222 (100%)	0	100	100
1	B	220/260 (85%)	220 (100%)	0	100	100
1	C	215/260 (83%)	214 (100%)	1 (0%)	88	92
1	D	218/260 (84%)	218 (100%)	0	100	100
1	E	218/260 (84%)	218 (100%)	0	100	100
1	F	218/260 (84%)	218 (100%)	0	100	100
1	G	217/260 (84%)	217 (100%)	0	100	100
1	H	217/260 (84%)	217 (100%)	0	100	100
2	a	198/194 (102%)	194 (98%)	4 (2%)	55	58
2	b	195/194 (100%)	193 (99%)	2 (1%)	76	81
2	c	195/194 (100%)	193 (99%)	2 (1%)	76	81
2	d	200/194 (103%)	196 (98%)	4 (2%)	55	58
2	e	196/194 (101%)	191 (97%)	5 (3%)	46	48
2	f	200/194 (103%)	197 (98%)	3 (2%)	65	69
2	g	196/194 (101%)	194 (99%)	2 (1%)	76	81
2	h	197/194 (102%)	195 (99%)	2 (1%)	76	81
All	All	3322/3632 (92%)	3297 (99%)	25 (1%)	84	86

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	d	121[A]	GLN
2	e	1	MSE
2	h	100	GLU
2	d	121[B]	GLN
2	e	7[A]	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/297 (83%)	-0.43	1 (0%) 92 92	14, 23, 38, 64	0
1	B	247/297 (83%)	-0.47	1 (0%) 92 92	14, 23, 37, 62	0
1	C	247/297 (83%)	-0.53	1 (0%) 92 92	11, 21, 36, 74	0
1	D	247/297 (83%)	-0.48	0 100 100	14, 24, 36, 61	0
1	E	247/297 (83%)	-0.44	2 (0%) 86 85	11, 23, 41, 77	0
1	F	247/297 (83%)	-0.47	1 (0%) 92 92	13, 23, 42, 72	0
1	G	248/297 (83%)	-0.40	1 (0%) 92 92	13, 22, 42, 89	0
1	H	247/297 (83%)	-0.37	1 (0%) 92 92	15, 26, 40, 67	0
2	a	225/231 (97%)	0.05	9 (4%) 38 37	17, 31, 60, 91	1 (0%)
2	b	225/231 (97%)	0.08	10 (4%) 34 33	15, 31, 62, 133	1 (0%)
2	c	222/231 (96%)	-0.01	9 (4%) 37 36	12, 29, 55, 110	0
2	d	225/231 (97%)	-0.11	4 (1%) 68 66	16, 28, 55, 75	0
2	e	224/231 (96%)	-0.22	5 (2%) 62 60	13, 24, 48, 79	0
2	f	223/231 (96%)	-0.30	5 (2%) 62 60	14, 23, 42, 83	0
2	g	222/231 (96%)	-0.14	3 (1%) 75 74	15, 27, 49, 80	0
2	h	226/231 (97%)	-0.16	5 (2%) 62 60	14, 26, 47, 62	1 (0%)
All	All	3769/4224 (89%)	-0.28	58 (1%) 73 72	11, 25, 47, 133	3 (0%)

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	a	226	THR	7.4
2	e	67	LEU	6.6
2	b	2	ALA	6.2
2	c	226	THR	5.7
2	a	227	PRO	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.