



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 27, 2024 – 04:27 pm BST

PDB ID : 3ZNS
Title : HDAC7 bound with TFMO inhibitor tmp942
Authors : Lobera, M.; Madauss, K.P.; Pohlhaus, D.T.; Trump, R.P.; Nolan, M.A.
Deposited on : 2013-02-15
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	423	
1	B	423	
1	C	423	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HISTONE DEACETYLASE 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	0	0
			2774	1740	499	516	19			
1	B	380	Total	C	N	O	S	0	0	0
			2867	1798	514	536	19			
1	C	359	Total	C	N	O	S	0	0	0
			2704	1691	490	505	18			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	GLY	-	expression tag	UNP Q8WUI4
B	481	GLY	-	expression tag	UNP Q8WUI4
C	481	GLY	-	expression tag	UNP Q8WUI4

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	K	0	0
			2	2		
3	B	2	Total	K	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	2	Total K 2 2	0	0

- Molecule 4 is N-{[1-methyl-4-(4-phenyl-1,3-thiazol-2-yl)piperidin-4-yl]methyl}-3-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide (three-letter code: NU7) (formula: C₂₆H₂₄F₃N₅O₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 37	C 26	F 3	N 5	O 2	S 1	0	0
4	B	1	Total 37	C 26	F 3	N 5	O 2	S 1	0	0
4	C	1	Total 37	C 26	F 3	N 5	O 2	S 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	46	Total O 46 46	0	0
5	B	69	Total O 69 69	0	0
5	C	18	Total O 18 18	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HISTONE DEACETYLASE 7

Chain A:




- Molecule 1: HISTONE DEACETYLASE 7

Chain B:



- Molecule 1: HISTONE DEACETYLASE 7

Chain C:



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	81.57Å 81.57Å 150.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	70.71 – 2.45 33.14 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.8 (70.71-2.45) 99.8 (33.14-2.45)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.88 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.5.0053	Depositor
R, R_{free}	0.186 , 0.225 0.239 , 0.271	Depositor DCC
R_{free} test set	2054 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	50.0	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.011 for -h,-k,l 0.045 for h,-h-k,-l 0.029 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8601	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NU7, ZN, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/2842	0.49	0/3860
1	B	0.33	0/2934	0.48	0/3981
1	C	0.31	0/2770	0.47	0/3764
All	All	0.32	0/8546	0.48	0/11605

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2774	0	2651	19	0
1	B	2867	0	2751	20	0
1	C	2704	0	2571	25	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
4	A	37	0	24	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	37	0	24	2	0
4	C	37	0	24	0	0
5	A	46	0	0	0	0
5	B	69	0	0	0	0
5	C	18	0	0	0	0
All	All	8601	0	8045	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:535:CYS:HG	2:C:102:ZN:ZN	0.69	0.94
1:C:623:VAL:HG22	1:C:627:THR:HB	1.66	0.77
1:B:623:VAL:HG22	1:B:627:THR:HB	1.67	0.76
1:B:523:ILE:HG21	1:B:644:SER:HB3	1.69	0.74
1:C:541:HIS:CE1	1:C:628:ILE:HD11	2.25	0.71
1:B:844:ASP:OD2	1:B:846:THR:HG22	1.92	0.68
1:A:623:VAL:HG22	1:A:627:THR:HB	1.76	0.65
1:B:808:ALA:HB3	1:B:809:PRO:HD3	1.80	0.63
1:C:796:VAL:HB	1:C:837:LEU:HD22	1.81	0.62
1:A:554:ARG:NH2	1:A:850:ASP:OD1	2.33	0.61
1:A:520:THR:HG23	1:A:860:LEU:HD23	1.82	0.61
1:C:651:LYS:HB3	1:C:657:LEU:HD13	1.84	0.60
1:C:592:ASN:CB	1:C:593:PRO:HA	2.33	0.59
1:A:744:VAL:HG12	1:A:890:VAL:HG21	1.85	0.58
1:A:649:ALA:HB1	1:A:692:LEU:HD21	1.86	0.57
1:C:688:ALA:O	1:C:692:LEU:HD13	2.05	0.57
1:C:548:ILE:HD13	1:C:665:ARG:HG2	1.88	0.55
1:B:796:VAL:HB	1:B:837:LEU:HD22	1.89	0.54
1:C:650:PHE:CE1	1:C:692:LEU:HD12	2.43	0.53
1:C:592:ASN:HB2	1:C:593:PRO:HA	1.91	0.53
1:B:520:THR:HG22	1:B:659:ASN:OD1	2.09	0.52
1:C:564:CYS:SG	1:C:860:LEU:HD21	2.49	0.52
1:B:810:LEU:HD22	4:B:1900:NU7:H29	1.92	0.52
1:A:519:THR:HG22	1:A:520:THR:H	1.74	0.52
1:B:661:PHE:CE2	1:B:663:VAL:HG22	2.46	0.50
1:B:649:ALA:HB1	1:B:692:LEU:HD21	1.92	0.49
1:C:806:HIS:O	1:C:812:GLY:N	2.38	0.49
1:C:796:VAL:CB	1:C:837:LEU:HD22	2.42	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:577:LEU:HD23	1:B:687:ILE:HD13	1.95	0.49
1:C:592:ASN:CB	1:C:593:PRO:CA	2.91	0.49
1:A:796:VAL:HB	1:A:837:LEU:HD22	1.94	0.48
1:A:556:GLN:HE22	1:B:579:SER:HA	1.78	0.48
1:A:795:LEU:HD23	1:A:836:VAL:HB	1.97	0.47
1:B:697:LYS:O	1:B:698:ALA:HB3	2.15	0.46
1:C:792:ASP:O	1:C:831:ALA:HB1	2.16	0.46
1:C:757:VAL:HG22	1:C:891:HIS:CE1	2.50	0.46
1:A:810:LEU:HD22	4:A:1900:NU7:C29	2.46	0.46
1:A:824:THR:CG2	1:A:859:LEU:HD13	2.46	0.45
1:A:520:THR:HB	1:A:659:ASN:OD1	2.16	0.45
1:C:824:THR:HG21	1:C:859:LEU:HD13	1.99	0.44
1:A:525:ASP:OD1	1:A:527:VAL:HG23	2.17	0.44
1:A:577:LEU:HD23	1:A:687:ILE:HD13	1.99	0.44
1:C:628:ILE:HG22	1:C:629:TRP:N	2.31	0.44
1:B:702:LEU:HD13	1:B:725:LEU:HD23	2.00	0.44
1:B:810:LEU:HD22	4:B:1900:NU7:C29	2.47	0.44
1:A:523:ILE:HG21	1:A:644:SER:HB3	1.99	0.44
1:A:817:ALA:HB2	1:A:850:ASP:HB3	2.00	0.43
1:B:548:ILE:HD13	1:B:665:ARG:HG2	2.01	0.43
1:C:592:ASN:HB3	1:C:593:PRO:HA	2.00	0.43
1:C:560:LEU:HD13	1:C:856:VAL:HG12	2.01	0.43
1:A:649:ALA:HA	1:A:836:VAL:HG21	2.01	0.43
1:B:731:ARG:CZ	1:B:765:LEU:HD11	2.49	0.42
1:C:554:ARG:NH2	1:C:850:ASP:OD1	2.53	0.42
1:C:529:LEU:HD23	1:C:548:ILE:HD11	2.01	0.41
1:B:592:ASN:O	1:B:595:SER:N	2.53	0.41
1:A:824:THR:HG21	1:A:859:LEU:HD13	2.02	0.41
1:C:592:ASN:HB3	1:C:593:PRO:CA	2.49	0.41
1:B:796:VAL:CB	1:B:837:LEU:HD22	2.51	0.40
1:A:757:VAL:HG22	1:A:891:HIS:CE1	2.56	0.40
1:B:757:VAL:HG22	1:B:891:HIS:CE1	2.56	0.40
1:B:796:VAL:CG2	1:B:837:LEU:HD22	2.51	0.40
1:C:824:THR:CG2	1:C:859:LEU:HD13	2.52	0.40
1:C:541:HIS:ND1	1:C:628:ILE:HD11	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/423 (86%)	347 (95%)	16 (4%)	1 (0%)	41	49
1	B	376/423 (89%)	363 (96%)	12 (3%)	1 (0%)	41	49
1	C	355/423 (84%)	341 (96%)	11 (3%)	3 (1%)	19	22
All	All	1095/1269 (86%)	1051 (96%)	39 (4%)	5 (0%)	29	34

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	592	ASN
1	A	811	GLY
1	C	811	GLY
1	B	811	GLY
1	C	673	HIS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/336 (86%)	277 (96%)	12 (4%)	30	39
1	B	298/336 (89%)	284 (95%)	14 (5%)	26	34
1	C	280/336 (83%)	276 (99%)	4 (1%)	67	77
All	All	867/1008 (86%)	837 (96%)	30 (4%)	36	47

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	519	THR
1	A	520	THR
1	A	532	GLN
1	A	543	GLU
1	A	553	SER
1	A	647	ASP
1	A	657	LEU
1	A	669	HIS
1	A	692	LEU
1	A	706	TRP
1	A	734	ASP
1	A	859	LEU
1	B	543	GLU
1	B	561	ARG
1	B	574	LEU
1	B	594	LEU
1	B	623	VAL
1	B	647	ASP
1	B	669	HIS
1	B	706	TRP
1	B	765	LEU
1	B	766	ASP
1	B	839	LEU
1	B	862	ASN
1	B	874	GLN
1	B	878	LEU
1	C	523	ILE
1	C	561	ARG
1	C	669	HIS
1	C	765	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	556	GLN
1	A	636	ASN
1	A	673	HIS
1	A	693	GLN
1	A	720	GLN
1	A	756	ASN
1	A	879	ASN
1	A	891	HIS
1	A	899	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	556	GLN
1	B	636	ASN
1	B	825	GLN
1	B	879	ASN
1	B	899	GLN
1	C	636	ASN
1	C	673	HIS
1	C	693	GLN
1	C	720	GLN
1	C	899	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 12 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NU7	A	1900	-	35,41,41	1.15	2 (5%)	48,60,60	1.88	8 (16%)
4	NU7	C	1900	2	35,41,41	1.13	2 (5%)	48,60,60	1.94	7 (14%)
4	NU7	B	1900	-	35,41,41	1.13	2 (5%)	48,60,60	1.93	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NU7	A	1900	-	-	0/23/42/42	0/5/5/5
4	NU7	C	1900	2	-	5/23/42/42	0/5/5/5
4	NU7	B	1900	-	-	7/23/42/42	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1900	NU7	C29-S30	4.29	1.77	1.70
4	C	1900	NU7	C29-S30	4.29	1.77	1.70
4	B	1900	NU7	C29-S30	3.94	1.76	1.70
4	B	1900	NU7	C07-N06	3.54	1.40	1.35
4	A	1900	NU7	C07-N06	3.47	1.40	1.35
4	C	1900	NU7	C07-N06	3.14	1.39	1.35

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1900	NU7	C10-C07-N08	6.68	127.81	119.12
4	B	1900	NU7	C29-C28-C31	-6.21	120.81	129.44
4	C	1900	NU7	C29-C28-C31	-5.31	122.05	129.44
4	C	1900	NU7	C07-N06-C05	5.26	108.84	101.38
4	B	1900	NU7	C28-C29-S30	-5.26	105.33	111.79
4	A	1900	NU7	C29-C28-C31	-5.25	122.14	129.44
4	B	1900	NU7	C10-C07-N08	5.16	125.83	119.12
4	C	1900	NU7	C28-C29-S30	-5.02	105.62	111.79
4	B	1900	NU7	C07-N06-C05	5.02	108.49	101.38
4	A	1900	NU7	C28-C29-S30	-4.99	105.66	111.79
4	A	1900	NU7	C10-C07-N08	4.83	125.41	119.12
4	A	1900	NU7	C07-N06-C05	4.76	108.12	101.38
4	A	1900	NU7	C02-C05-N06	4.43	126.73	122.41
4	C	1900	NU7	C20-C19-N18	-3.57	109.17	114.12
4	B	1900	NU7	C02-C05-N06	3.51	125.83	122.41
4	C	1900	NU7	C21-C22-N2	3.15	113.18	110.82
4	A	1900	NU7	C20-C19-N18	-3.02	109.94	114.12
4	A	1900	NU7	F04-C02-C05	-2.96	106.17	112.24
4	B	1900	NU7	F04-C02-C05	-2.86	106.37	112.24
4	B	1900	NU7	C21-C22-N2	2.72	112.86	110.82
4	C	1900	NU7	F03-C02-C05	-2.45	107.21	112.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1900	NU7	C21-C22-N2	2.24	112.50	110.82

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1900	NU7	C13-C14-C16-N18
4	B	1900	NU7	C15-C14-C16-N18
4	B	1900	NU7	C13-C14-C16-O17
4	B	1900	NU7	C15-C14-C16-O17
4	C	1900	NU7	C13-C14-C16-N18
4	C	1900	NU7	C13-C14-C16-O17
4	C	1900	NU7	C15-C14-C16-N18
4	C	1900	NU7	C20-C19-N18-C16
4	C	1900	NU7	C15-C14-C16-O17
4	B	1900	NU7	F04-C02-C05-N06
4	B	1900	NU7	F01-C02-C05-N06
4	B	1900	NU7	F03-C02-C05-N06

There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1900	NU7	1	0
4	B	1900	NU7	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand NU7 C 1900




Bond lengths



Bond angles




Torsions




Rings


Ligand NU7 B 1900




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NU7 B 1900:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NU7 C 1900:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.