wwPDB X-ray Structure Validation Summary Report (i)

May 4, 2024 – 09:42 am BST

PDB ID	:	7ZOO
Title	:	Carbohydrate binding domain CBM92-B from a multi-catalytic glucanase-chitinase from Chitinophaga pinensis DSM 2588 in complex with gentiobiose
Authors	:	Mazurkewich, S.; McKee, L.S.; Lu, Z.; Branden, G.; Larsbrink, J.
Deposited on	:	2022-04-26
Resolution	:	1.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	147	3% 80% 5% 15%
1	B	147	4% 80% 5% 14%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoside hydrolase family 18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	125	964	596	171	191	6	0	0	0
1	B	126	970	599	172	193	6	0	0	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	874	MET	-	initiating methionine	UNP A0A979GQH9
A	875	ALA	-	expression tag	UNP A0A979GQH9
A	876	SER	-	expression tag	UNP A0A979GQH9
A	877	MET	-	expression tag	UNP A0A979GQH9
A	878	THR	-	expression tag	UNP A0A979GQH9
A	879	GLY	-	expression tag	UNP A0A979GQH9
A	880	GLY	-	expression tag	UNP A0A979GQH9
A	881	GLN	-	expression tag	UNP A0A979GQH9
A	882	GLN	-	expression tag	UNP A0A979GQH9
A	883	MET	-	expression tag	UNP A0A979GQH9
A	884	GLY	-	expression tag	UNP A0A979GQH9
A	885	ARG	-	expression tag	UNP A0A979GQH9
A	886	GLY	-	expression tag	UNP A0A979GQH9
A	887	SER	-	expression tag	UNP A0A979GQH9
A	959	SNN	ASP	conflict	UNP A0A979GQH9
A	1013	LEU	-	expression tag	UNP A0A979GQH9
A	1014	GLU	-	expression tag	UNP A0A979GQH9
A	1015	HIS	-	expression tag	UNP A0A979GQH9
A	1016	HIS	-	expression tag	UNP A0A979GQH9
A	1017	HIS	-	expression tag	UNP A0A979GQH9
A	1018	HIS	-	expression tag	UNP A0A979GQH9
A	1019	HIS	-	expression tag	UNP A0A979GQH9
A	1020	HIS	-	expression tag	UNP A0A979GQH9
B	874	MET	-	initiating methionine	UNP A0A979GQH9
B	875	ALA	-	expression tag	UNP A0A979GQH9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	876	SER	-	expression tag	UNP A0A979GQH9
B	877	MET	-	expression tag	UNP A0A979GQH9
B	878	THR	-	expression tag	UNP A0A979GQH9
B	879	GLY	-	expression tag	UNP A0A979GQH9
B	880	GLY	-	expression tag	UNP A0A979GQH9
B	881	GLN	-	expression tag	UNP A0A979GQH9
B	882	GLN	-	expression tag	UNP A0A979GQH9
B	883	MET	-	expression tag	UNP A0A979GQH9
B	884	GLY	-	expression tag	UNP A0A979GQH9
B	885	ARG	-	expression tag	UNP A0A979GQH9
B	886	GLY	-	expression tag	UNP A0A979GQH9
B	887	SER	-	expression tag	UNP A0A979GQH9
B	959	SNN	ASP	conflict	UNP A0A979GQH9
B	1013	LEU	-	expression tag	UNP A0A979GQH9
B	1014	GLU	-	expression tag	UNP A0A979GQH9
B	1015	HIS	-	expression tag	UNP A0A979GQH9
B	1016	HIS	-	expression tag	UNP A0A979GQH9
B	1017	HIS	-	expression tag	UNP A0A979GQH9
B	1018	HIS	-	expression tag	UNP A0A979GQH9
B	1019	HIS	-	expression tag	UNP A0A979GQH9
B	1020	HIS	-	expression tag	UNP A0A979GQH9

- Molecule 2 is beta-D-glucopyranose (three-letter code: BGC) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	12	6	6	0	0
			Total	C	O		
2	A	1	12	6	6	0	0
			Total	C	O		
2	B	1	12	6	6	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	69	69	69	0	0
			Total	O		
3	B	59	59	59	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycoside hydrolase family 18

Chain A:



- Molecule 1: Glycoside hydrolase family 18

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	44.62Å 111.41Å 70.54Å 90.00° 98.58° 90.00°	Depositor
Resolution (Å)	29.56 – 1.84 29.56 – 1.84	Depositor EDS
% Data completeness (in resolution range)	95.5 (29.56-1.84) 95.5 (29.56-1.84)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.32 (at 1.84Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.191 , 0.216 0.192 , 0.216	Depositor DCC
R_{free} test set	2098 reflections (7.50%)	wwPDB-VP
Wilson B-factor (Å^2)	35.2	Xtriage
Anisotropy	0.736	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 45.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	2098	wwPDB-VP
Average B, all atoms (Å^2)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, SNN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.37	0/978	0.54	0/1327
1	B	0.38	0/984	0.54	0/1335
All	All	0.38	0/1962	0.54	0/2662

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	958	ILE	Peptide
1	B	958	ILE	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	964	0	878	4	0
1	B	970	0	883	3	0
2	A	24	0	24	0	0
2	B	12	0	12	0	0
3	A	69	0	0	1	0
3	B	59	0	0	0	0
All	All	2098	0	1797	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 7 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:946:ASN:HB2	1:A:961:TRP:CE2	2.49	0.46
1:A:888:LYS:HB2	1:A:888:LYS:HE2	1.60	0.46
1:B:899:TYR:CE1	1:B:916:PRO:HD3	2.53	0.43
1:A:965:ASP:HB2	1:A:977:ARG:HB3	2.02	0.41
1:B:940:LEU:HD22	1:B:956:PRO:HA	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	121/147 (82%)	114 (94%)	7 (6%)	0	100 100
1	B	122/147 (83%)	112 (92%)	10 (8%)	0	100 100
All	All	243/294 (83%)	226 (93%)	17 (7%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/116 (85%)	99 (100%)	0	100	100
1	B	100/116 (86%)	100 (100%)	0	100	100
All	All	199/232 (86%)	199 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	SNN	B	959	1	7,8,8	1.19	1 (14%)	7,11,11	3.53	5 (71%)
1	SNN	A	959	1	7,8,8	1.53	2 (28%)	7,11,11	2.95	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SNN	B	959	1	-	-	0/1/1/1
1	SNN	A	959	1	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	959	SNN	C-N1	-2.94	1.33	1.37
1	A	959	SNN	C5-N1	-2.56	1.33	1.37
1	B	959	SNN	C-N1	-2.49	1.34	1.37

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	959	SNN	O-C-CA	-5.83	121.96	126.18
1	A	959	SNN	CA-C-N1	5.26	111.36	107.30
1	B	959	SNN	CA-C-N1	4.65	110.89	107.30
1	A	959	SNN	O-C-CA	-4.50	122.93	126.18
1	B	959	SNN	O5-C5-C4	-3.98	121.14	126.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	BGC	A	1102	-	12,12,12	0.55	0	17,17,17	0.77	0
2	BGC	A	1101	-	12,12,12	0.46	0	17,17,17	1.03	1 (5%)
2	BGC	B	1101	-	12,12,12	0.50	0	17,17,17	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	1102	-	-	0/2/22/22	0/1/1/1
2	BGC	A	1101	-	-	0/2/22/22	0/1/1/1
2	BGC	B	1101	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	1101	BGC	C6-C5-C4	-2.14	108.00	113.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand BGC A 1102



Bond lengths



Bond angles



Torsions



Rings

Ligand BGC A 1101



Ligand BGC B 1101



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	124/147 (84%)	0.14	5 (4%)	38	35	32, 40, 52, 72	0
1	B	125/147 (85%)	0.09	6 (4%)	30	28	31, 40, 53, 59	0
All	All	249/294 (84%)	0.12	11 (4%)	34	31	31, 40, 53, 72	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	895	PHE	5.8
1	A	1012	THR	4.2
1	A	970	SER	3.5
1	A	1011	ALA	3.3
1	B	887	SER	3.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	SNN	A	959	8/8	0.93	0.14	37,40,47,51	0
1	SNN	B	959	8/8	0.94	0.09	41,44,47,47	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	BGC	A	1102	12/12	0.92	0.13	39,45,48,55	0
2	BGC	B	1101	12/12	0.93	0.10	43,48,51,58	0
2	BGC	A	1101	12/12	0.94	0.10	45,48,54,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BGC A 1102:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BGC B 1101:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BGC A 1101:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.